

Product datasheet for **SR401799**

Egln3 Mouse siRNA Oligo Duplex (Locus ID 112407)

Product data:

Product Type:	siRNA Oligo Duplexes
Purity:	HPLC purified
Quality Control:	Tested by ESI-MS
Sequences:	Available with shipment
Stability:	One year from date of shipment when stored at -20°C.
# of transfections:	Approximately 330 transfections/2nmol in 24-well plate under optimized conditions (final conc. 10 nM).
Note:	Single siRNA duplex (10nmol) can be ordered.
RefSeq:	NM_028133
UniProt ID:	Q91UZ4
Synonyms:	2610021G09Rik; AI505553; AI648162; Hif-p4h-3; Phd3; SM-20
Components:	Egln3 (Mouse) - 3 unique 27mer siRNA duplexes - 2 nmol each (Locus ID 112407) Included - SR30004, Trilencer-27 Universal Scrambled Negative Control siRNA Duplex - 2 nmol Included - SR30005, RNase free siRNA Duplex Resuspension Buffer - 2 ml



[View online »](#)

Summary:

Plays a crucial role in DNA damage response (DDR) by hydroxylating TELO2, promoting its interaction with ATR which is required for activation of the ATR/CHK1/p53 pathway (By similarity). Cellular oxygen sensor that catalyzes, under normoxic conditions, the post-translational formation of 4-hydroxyproline in hypoxia-inducible factor (HIF) alpha proteins. Hydroxylates a specific proline found in each of the oxygen-dependent degradation (ODD) domains (N-terminal, NODD, and C-terminal, CODD) of HIF1A. Also hydroxylates HIF2A. Has a preference for the CODD site for both HIF1A and HIF2A. Hydroxylation on the NODD site by EGLN3 appears to require prior hydroxylation on the CODD site. Hydroxylated HIFs are then targeted for proteasomal degradation via the von Hippel-Lindau ubiquitination complex. Under hypoxic conditions, the hydroxylation reaction is attenuated allowing HIFs to escape degradation resulting in their translocation to the nucleus, heterodimerization with HIF1B, and increased expression of hypoxia-inducible genes. EGLN3 is the most important isozyme in limiting physiological activation of HIFs (particularly HIF2A) in hypoxia. Also hydroxylates PKM in hypoxia, limiting glycolysis. Under normoxia, hydroxylates and regulates the stability of ADRB2. Regulator of cardiomyocyte and neuronal apoptosis. In cardiomyocytes, inhibits the anti-apoptotic effect of BCL2 by disrupting the BAX-BCL2 complex. In neurons, has a NGF-induced proapoptotic effect, probably through regulating CASP3 activity. Also essential for hypoxic regulation of neutrophilic inflammation. Target proteins are preferentially recognized via a LXXLAP motif.[UniProtKB/Swiss-Prot Function]

Performance Guaranteed:

OriGene guarantees that at least two of the three Dicer-Substrate duplexes in the kit will provide at least 70% or more knockdown of the target mRNA when used at 10 nM concentration by quantitative RT-PCR when the TYE-563 fluorescent transfection control duplex (cat# SR30002) indicates that >90% of the cells have been transfected and the HPRT positive control (cat# SR30003) provides 90% knockdown efficiency.

For non-conforming siRNA, requests for replacement product must be made within ninety (90) days from the date of delivery of the siRNA kit. To arrange for a free replacement with newly designed duplexes, please contact Technical Services at techsupport@origene.com. Please provide your data indicating the transfection efficiency and measurement of gene expression knockdown compared to the scrambled siRNA control (quantitative RT-PCR data required).