

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for SR405143

Tmed2 Mouse siRNA Oligo Duplex (Locus ID 56334)

Product data:

| Product Type: | siRNA Oligo Duplexes |
|---------------------|---|
| Purity: | HPLC purified |
| Quality Control: | Tested by ESI-MS |
| Sequences: | Available with shipment |
| Stability: | One year from date of shipment when stored at -20°C. |
| # of transfections: | Approximately 330 transfections/2nmol in 24-well plate under optimized conditions (final conc. 10 nM). |
| Note: | Single siRNA duplex (10nmol) can be ordered. |
| RefSeq: | <u>NM 019770</u> |
| UniProt ID: | <u>Q9R0Q3</u> |
| Synonyms: | 1110032D12Rik; 1810020N21Rik; p24beta1; Rnp24; Sid394 |
| Components: | Tmed2 (Mouse) - 3 unique 27mer siRNA duplexes - 2 nmol each (Locus ID 56334) Included - SR30004, Trilencer-27 Universal Scrambled Negative Control siRNA Duplex - 2 nmol Included - SR30005, RNAse free siRNA Duplex Resuspension Buffer - 2 ml |



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2021 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

GRIGENE Tmed2 Mouse siRNA Oligo Duplex (Locus ID 56334) – SR405143

Involved in vesicular protein trafficking. Mainly functions in the early secretory pathway but Summary: also in post-Golgi membranes. Thought to act as cargo receptor at the lumenal side for incorporation of secretory cargo molecules into transport vesicles and to be involved in vesicle coat formation at the cytoplasmic side. In COPII vesicle-mediated anterograde transport involved in the transport of GPI-anchored proteins and proposed to act together with TMED10 as their cargo receptor; the function specifically implies SEC24C and SEC24D of the COPII vesicle coat and lipid raft-like microdomains of the ER. Recognizes GPI anchors structural remodeled in the ER by PGAP1 and MPPE1. In COPI vesicle-mediated retrograde transport inhibits the GTPase-activating activity of ARFGAP1 towards ARF1 thus preventing immature uncoating and allowing cargo selection to take place. Involved in trafficking of G protein-coupled receptors (GPCRs). Regulates F2RL1, OPRM1 and P2RY4 exocytic trafficking from the Golgi to the plasma membrane thus contributing to receptor resensitization. Facilitates CASR maturation and stabilization in the early secretory pathway and increases CASR plasma membrane targeting. Proposed to be involved in organization of intracellular membranes such as the maintenance of the Golgi apparatus. May also play a role in the biosynthesis of secreted cargo such as eventual processing (By similarity). Required for morphogenesis of embryo and placenta.[UniProtKB/Swiss-Prot Function] Performance OriGene guarantees that at least two of the three Dicer-Substrate duplexes in the kit will **Guaranteed:** provide at least 70% or more knockdown of the target mRNA when used at 10 nM

positive control (cat# SR30003) provides 90% knockdown efficiency.

concentration by quantitative RT-PCR when the TYE-563 fluorescent transfection control duplex (cat# SR30002) indicates that >90% of the cells have been transfected and the HPRT

For non-conforming siRNA, requests for replacement product must be made within ninety (90) days from the date of delivery of the siRNA kit. To arrange for a free replacement with newly designed duplexes, please contact Technical Services at techsupport@origene.com. Please provide your data indicating the transfection efficiency and measurement of gene expression knockdown compared to the scrambled siRNA control (quantitative RT-PCR data

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2021 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

required).