

Product datasheet for SR405655

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Otulin Mouse siRNA Oligo Duplex (Locus ID 432940)

Product data:

Product Type: siRNA Oligo Duplexes

Purity: HPLC purified

Quality Control: Tested by ESI-MS

Sequences: Available with shipment

Stability: One year from date of shipment when stored at -20°C.

of transfections: Approximately 330 transfections/2nmol in 24-well plate under optimized conditions (final

conc. 10 nM).

Note: Single siRNA duplex (10nmol) can be ordered.

RefSeq: <u>NM 001013792</u>

UniProt ID: Q3UCV8

Synonyms: C79097; Fam105b; m3Sapc; m7-1Sapc

Components: Otulin (Mouse) - 3 unique 27mer siRNA duplexes - 2 nmol each (Locus ID 432940)

Included - SR30004, Trilencer-27 Universal Scrambled Negative Control siRNA Duplex - 2 nmol

Included - SR30005, RNAse free siRNA Duplex Resuspension Buffer - 2 ml

Summary: Deubiquitinase that specifically removes linear ('Met-1'-linked) polyubiquitin chains to

substrates and acts as a regulator of angiogenesis and innate immune response

(PubMed:23708998, PubMed:27523608, PubMed:29950720). Required during angiogenesis, craniofacial and neuronal development by regulating the canonical Wnt signaling together with the LUBAC complex (PubMed:23708998). Acts as a negative regulator of NF-kappa-B by regulating the activity of the LUBAC complex (By similarity). OTULIN function is mainly restricted to homeostasis of the LUBAC complex: acts by removing 'Met-1'-linked

autoubiquitination of the LUBAC complex, thereby preventing inactivation of the LUBAC complex (PubMed:29950720). Acts as a key negative regulator of inflammation by restricting spontaneous inflammation and maintaining immune homeostasis (PubMed:27523608, PubMed:29950720). In myeloid cell, required to prevent unwarranted secretion of cytokines leading to inflammation and autoimmunity by restricting linear polyubiquitin formation

(PubMed:27523608). Plays a role in innate immune response by restricting linear

polyubiquitin formation on LUBAC complex in response to NOD2 stimulation, probably to limit NOD2-dependent proinflammatory signaling (By similarity).[UniProtKB/Swiss-Prot

Function]







Performance Guaranteed:

OriGene guarantees that at least two of the three Dicer-Substrate duplexes in the kit will provide at least 70% or more knockdown of the target mRNA when used at 10 nM concentration by quantitative RT-PCR when the TYE-563 fluorescent transfection control duplex (cat# SR30002) indicates that >90% of the cells have been transfected and the HPRT positive control (cat# SR30003) provides 90% knockdown efficiency.

For non-conforming siRNA, requests for replacement product must be made within ninety (90) days from the date of delivery of the siRNA kit. To arrange for a free replacement with newly designed duplexes, please contact Technical Services at techsupport@origene.com. Please provide your data indicating the transfection efficiency and measurement of gene expression knockdown compared to the scrambled siRNA control (quantitative RT-PCR data required).