

Product datasheet for SR408383

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Exosc7 Mouse siRNA Oligo Duplex (Locus ID 66446)

Product data:

Product Type: siRNA Oligo Duplexes

Purity: HPLC purified

Quality Control: Tested by ESI-MS

Sequences: Available with shipment

Stability: One year from date of shipment when stored at -20°C.

of transfections: Approximately 330 transfections/2nmol in 24-well plate under optimized conditions (final

conc. 10 nM).

Note: Single siRNA duplex (10nmol) can be ordered.

RefSeq: <u>NM 001081188</u>

UniProt ID: Q9D0M0

Synonyms: 2610002K22Rik; AV212732; mKIAA0116

Components: Exosc7 (Mouse) - 3 unique 27mer siRNA duplexes - 2 nmol each (Locus ID 66446)

Included - SR30004, Trilencer-27 Universal Scrambled Negative Control siRNA Duplex - 2 nmol

Included - SR30005, RNAse free siRNA Duplex Resuspension Buffer - 2 ml

Summary: Non-catalytic component of the RNA exosome complex which has 3'->5' exoribonuclease

activity and participates in a multitude of cellular RNA processing and degradation events. In the nucleus, the RNA exosome complex is involved in proper maturation of stable RNA species such as rRNA, snRNA and snoRNA, in the elimination of RNA processing by-products and non-coding 'pervasive' transcripts, such as antisense RNA species and promoter-

upstream transcripts (PROMPTs), and of mRNAs with processing defects, thereby limiting or excluding their export to the cytoplasm. The RNA exosome may be involved in Ig class switch recombination (CSR) and/or Ig variable region somatic hypermutation (SHM) by targeting

AICDA deamination activity to transcribed dsDNA substrates. In the cytoplasm, the RNA exosome complex is involved in general mRNA turnover and specifically degrades inherently unstable mRNAs containing AU-rich elements (AREs) within their 3' untranslated regions, and in RNA surveillance pathways, preventing translation of aberrant mRNAs. It seems to be involved in degradation of histone mRNA. The catalytic inactive RNA exosome core complex of 9 subunits (Exo-9) is proposed to play a pivotal role in the binding and presentation of RNA for ribonucleolysis, and to serve as a scaffold for the association with catalytic subunits and

accessory proteins or complexes (By similarity).[UniProtKB/Swiss-Prot Function]







Performance Guaranteed:

OriGene guarantees that at least two of the three Dicer-Substrate duplexes in the kit will provide at least 70% or more knockdown of the target mRNA when used at 10 nM concentration by quantitative RT-PCR when the TYE-563 fluorescent transfection control duplex (cat# SR30002) indicates that >90% of the cells have been transfected and the HPRT positive control (cat# SR30003) provides 90% knockdown efficiency.

For non-conforming siRNA, requests for replacement product must be made within ninety (90) days from the date of delivery of the siRNA kit. To arrange for a free replacement with newly designed duplexes, please contact Technical Services at techsupport@origene.com. Please provide your data indicating the transfection efficiency and measurement of gene expression knockdown compared to the scrambled siRNA control (quantitative RT-PCR data required).