

Product datasheet for **SR411858**

Slamf6 Mouse siRNA Oligo Duplex (Locus ID 30925)

Product data:

Product Type:	siRNA Oligo Duplexes
Purity:	HPLC purified
Quality Control:	Tested by ESI-MS
Sequences:	Available with shipment
Stability:	One year from date of shipment when stored at -20°C.
# of transfections:	Approximately 330 transfections/2nmol in 24-well plate under optimized conditions (final conc. 10 nM).
Note:	Single siRNA duplex (10nmol) can be ordered.
RefSeq:	NM_001347186 , NM_001347187 , NM_030710
UniProt ID:	Q9ET39
Synonyms:	KAL1; KAL1b; Ly108; NTB-A; NTBA; SF2000
Components:	Slamf6 (Mouse) - 3 unique 27mer siRNA duplexes - 2 nmol each (Locus ID 30925) Included - SR30004, Trilencer-27 Universal Scrambled Negative Control siRNA Duplex - 2 nmol Included - SR30005, RNase free siRNA Duplex Resuspension Buffer - 2 ml



[View online »](#)

Summary:

Self-ligand receptor of the signaling lymphocytic activation molecule (SLAM) family. SLAM receptors triggered by homo- or heterotypic cell-cell interactions are modulating the activation and differentiation of a wide variety of immune cells and thus are involved in the regulation and interconnection of both innate and adaptive immune response. Activities are controlled by presence or absence of small cytoplasmic adapter proteins, SH2D1A/SAP and/or SH2D1B/EAT-2 (PubMed:19648922). Triggers cytolytic activity only in natural killer cells (NK) expressing high surface densities of natural cytotoxicity receptors (By similarity). Positive signaling in NK cells implicates phosphorylation of VAV1. NK cell activation seems to depend on SH2D1B and not on SH2D1A (By similarity). In conjunction with SLAMF1 controls the transition between positive selection and the subsequent expansion and differentiation of the thymocytic natural killer T (NKT) cell lineage (PubMed:18031695). Promotes T cell differentiation into a helper T-cell Th17 phenotype leading to increased IL-17 secretion; the costimulatory activity requires SH2D1A (By similarity). Promotes recruitment of RORC to the IL-17 promoter (By similarity). In conjunction with SLAMF1 and CD84/SLAMF5 may be a negative regulator of the humoral immune response (PubMed:25926831). In the absence of SH2D1A/SAP can transmit negative signals to CD4(+) T-cells and NKT cells. Negatively regulates germinal center formation by inhibiting T-cell:B-cell adhesion; the function probably implicates increased association with PTPN6/SHP-1 via ITSMs in absence of SH2D1A/SAP (PubMed:22683125). However, reported to mediated T-cell adhesion, to participate in stable T-cell:B-cell interactions and to be involved in maintaining B-cell tolerance in germinal centers and in preventing autoimmunity (PubMed:20153220, PubMed:25801429). Involved in regulation of autoimmunity. Isoform 3 may be suppressor of pathogenic T-cell proliferation (PubMed:21422172).[UniProtKB/Swiss-Prot Function]

Performance Guaranteed:

OriGene guarantees that at least two of the three Dicer-Substrate duplexes in the kit will provide at least 70% or more knockdown of the target mRNA when used at 10 nM concentration by quantitative RT-PCR when the TYE-563 fluorescent transfection control duplex (cat# SR30002) indicates that >90% of the cells have been transfected and the HPRT positive control (cat# SR30003) provides 90% knockdown efficiency.

For non-conforming siRNA, requests for replacement product must be made within ninety (90) days from the date of delivery of the siRNA kit. To arrange for a free replacement with newly designed duplexes, please contact Technical Services at techsupport@origene.com. Please provide your data indicating the transfection efficiency and measurement of gene expression knockdown compared to the scrambled siRNA control (quantitative RT-PCR data required).