

## Product datasheet for **SR416135**

### Fto Mouse siRNA Oligo Duplex (Locus ID 26383)

#### Product data:

Product Type:	siRNA Oligo Duplexes
Purity:	HPLC purified
Quality Control:	Tested by ESI-MS
Sequences:	Available with shipment
Stability:	One year from date of shipment when stored at -20°C.
# of transfections:	Approximately 330 transfections/2nmol in 24-well plate under optimized conditions (final conc. 10 nM).
Note:	Single siRNA duplex (10nmol) can be ordered.
RefSeq:	<a href="#">NM_011936</a>
UniProt ID:	<a href="#">Q8BGW1</a>
Synonyms:	AW743446; mKIAA1752
Components:	Fto (Mouse) - 3 unique 27mer siRNA duplexes - 2 nmol each (Locus ID 26383) Included - SR30004, Trilencer-27 Universal Scrambled Negative Control siRNA Duplex - 2 nmol Included - SR30005, RNase free siRNA Duplex Resuspension Buffer - 2 ml



[View online »](#)

**Summary:**

RNA demethylase that mediates oxidative demethylation of different RNA species, such as mRNAs, tRNAs and snRNAs, and acts as a regulator of fat mass, adipogenesis and energy homeostasis (PubMed:17991826, PubMed:18775698, PubMed:28002401). Specifically demethylates N(6)-methyladenosine (m6A) RNA, the most prevalent internal modification of messenger RNA (mRNA) in higher eukaryotes (PubMed:28002401). M6A demethylation by FTO affects mRNA expression and stability (By similarity). Also able to demethylate m6A in U6 small nuclear RNA (snRNA) (By similarity). Mediates demethylation of N(6),2'-O-dimethyladenosine cap (m6A(m)), by demethylating the N(6)-methyladenosine at the second transcribed position of mRNAs and U6 snRNA (PubMed:28002401). Demethylation of m6A(m) in the 5'-cap by FTO affects mRNA stability by promoting susceptibility to decapping (By similarity). Also acts as a tRNA demethylase by removing N(1)-methyladenine from various tRNAs (By similarity). Has no activity towards 1-methylguanine (By similarity). Has no detectable activity towards double-stranded DNA (By similarity). Also able to repair alkylated DNA and RNA by oxidative demethylation: demethylates single-stranded RNA containing 3-methyluracil, single-stranded DNA containing 3-methylthymine and has low demethylase activity towards single-stranded DNA containing 1-methyladenine or 3-methylcytosine (PubMed:17991826, PubMed:18775698). Ability to repair alkylated DNA and RNA is however unsure in vivo (PubMed:17991826, PubMed:18775698). Involved in the regulation of fat mass, adipogenesis and body weight, thereby contributing to the regulation of body size and body fat accumulation (PubMed:19234441, PubMed:19680540, PubMed:21076408, PubMed:23817550, PubMed:23300482). Involved in the regulation of thermogenesis and the control of adipocyte differentiation into brown or white fat cells (PubMed:19234441, PubMed:19680540). Regulates activity of the dopaminergic midbrain circuitry via its ability to demethylate m6A in mRNAs (PubMed:23817550).[UniProtKB/Swiss-Prot Function]

**Performance Guaranteed:**

OriGene guarantees that at least two of the three Dicer-Substrate duplexes in the kit will provide at least 70% or more knockdown of the target mRNA when used at 10 nM concentration by quantitative RT-PCR when the TYE-563 fluorescent transfection control duplex (cat# SR30002) indicates that >90% of the cells have been transfected and the HPRT positive control (cat# SR30003) provides 90% knockdown efficiency.

For non-conforming siRNA, requests for replacement product must be made within ninety (90) days from the date of delivery of the siRNA kit. To arrange for a free replacement with newly designed duplexes, please contact Technical Services at techsupport@origene.com. Please provide your data indicating the transfection efficiency and measurement of gene expression knockdown compared to the scrambled siRNA control (quantitative RT-PCR data required).