

## Product datasheet for **SR417439**

### **Fbxw11 Mouse siRNA Oligo Duplex (Locus ID 103583)**

#### **Product data:**

<b>Product Type:</b>	siRNA Oligo Duplexes
<b>Purity:</b>	HPLC purified
<b>Quality Control:</b>	Tested by ESI-MS
<b>Sequences:</b>	Available with shipment
<b>Stability:</b>	One year from date of shipment when stored at -20°C.
<b># of transfections:</b>	Approximately 330 transfections/2nmol in 24-well plate under optimized conditions (final conc. 10 nM).
<b>Note:</b>	Single siRNA duplex (10nmol) can be ordered.
<b>RefSeq:</b>	<a href="#">NM_001271347</a> , <a href="#">NM_001271348</a> , <a href="#">NM_001271349</a> , <a href="#">NM_134015</a> , <a href="#">NM_001363353</a> , <a href="#">NM_001363354</a> , <a href="#">NM_001363355</a>
<b>UniProt ID:</b>	<a href="#">Q5SRY7</a>
<b>Synonyms:</b>	2310065A07Rik; AA536858; BTRC2; BTRCP2; Fbxw1b; HOS
<b>Components:</b>	Fbxw11 (Mouse) - 3 unique 27mer siRNA duplexes - 2 nmol each (Locus ID 103583) Included - SR30004, Trilencer-27 Universal Scrambled Negative Control siRNA Duplex - 2 nmol Included - SR30005, RNase free siRNA Duplex Resuspension Buffer - 2 ml
<b>Summary:</b>	Substrate recognition component of a SCF (SKP1-CUL1-F-box protein) E3 ubiquitin-protein ligase complex which mediates the ubiquitination and subsequent proteasomal degradation of target proteins. Probably recognizes and binds to phosphorylated target proteins. SCF(FBXW11) mediates the ubiquitination of phosphorylated CTNNB1 and participates in Wnt signaling. SCF(FBXW11) mediates the ubiquitination of phosphorylated NFKBIA, which degradation frees the associated NFKB1 to translocate into the nucleus and to activate transcription. SCF(FBXW11) mediates the ubiquitination of IFNAR1. SCF(FBXW11) mediates the ubiquitination of CEP68; this is required for centriole separation during mitosis (By similarity). Involved in the oxidative stress-induced a ubiquitin-mediated decrease in RCAN1. Mediates the degradation of CDC25A induced by ionizing radiation in cells progressing through S phase and thus may function in the intra-S-phase checkpoint. Has an essential role in the control of the clock-dependent transcription via degradation of phosphorylated PER1 and phosphorylated PER2. SCF(FBXW11) mediates the ubiquitination of CYTH1, and probably CYTH2 (By similarity).[UniProtKB/Swiss-Prot Function]



[View online »](#)

**Performance  
Guaranteed:**

OriGene guarantees that at least two of the three Dicer-Substrate duplexes in the kit will provide at least 70% or more knockdown of the target mRNA when used at 10 nM concentration by quantitative RT-PCR when the TYE-563 fluorescent transfection control duplex (cat# SR30002) indicates that >90% of the cells have been transfected and the HPRT positive control (cat# SR30003) provides 90% knockdown efficiency.

For non-conforming siRNA, requests for replacement product must be made within ninety (90) days from the date of delivery of the siRNA kit. To arrange for a free replacement with newly designed duplexes, please contact Technical Services at [techsupport@origene.com](mailto:techsupport@origene.com). Please provide your data indicating the transfection efficiency and measurement of gene expression knockdown compared to the scrambled siRNA control (quantitative RT-PCR data required).