

## Product datasheet for **SR418390**

### Samhd1 Mouse siRNA Oligo Duplex (Locus ID 56045)

#### Product data:

Product Type:	siRNA Oligo Duplexes
Purity:	HPLC purified
Quality Control:	Tested by ESI-MS
Sequences:	Available with shipment
Stability:	One year from date of shipment when stored at -20°C.
# of transfections:	Approximately 330 transfections/2nmol in 24-well plate under optimized conditions (final conc. 10 nM).
Note:	Single siRNA duplex (10nmol) can be ordered.
RefSeq:	<a href="#">NM_001139520</a> , <a href="#">NM_018851</a> , <a href="#">NM_001370610</a>
UniProt ID:	<a href="#">Q60710</a>
Synonyms:	E330031J07Rik; Mg11
Components:	Samhd1 (Mouse) - 3 unique 27mer siRNA duplexes - 2 nmol each (Locus ID 56045) Included - SR30004, Trilencer-27 Universal Scrambled Negative Control siRNA Duplex - 2 nmol Included - SR30005, RNase free siRNA Duplex Resuspension Buffer - 2 ml



[View online »](#)

**Summary:**

Isoform 1: Protein that acts both as a host restriction factor involved in defense response to virus and as a regulator of DNA end resection at stalled replication forks (By similarity). Has deoxynucleoside triphosphate (dNTPase) activity, which is required to restrict infection by viruses: dNTPase activity reduces cellular dNTP levels to levels too low for retroviral reverse transcription to occur, blocking early-stage virus replication in dendritic and other myeloid cells (PubMed:23972988, PubMed:23872947, PubMed:26667483, PubMed:29379009). Likewise, suppresses LINE-1 retrotransposon activity (PubMed:26667483). In addition to virus restriction, dNTPase activity acts as a regulator of DNA precursor pools by regulating dNTP pools (By similarity). Phosphorylation at Thr-634 acts as a switch to control dNTPase-dependent and -independent functions: it inhibits dNTPase activity and ability to restrict infection by viruses, while it promotes DNA end resection at stalled replication forks (By similarity). Functions during S phase at stalled DNA replication forks to promote the resection of gapped or reversed forks: acts by stimulating the exonuclease activity of MRE11, activating the ATR-CHEK1 pathway and allowing the forks to restart replication (By similarity). Its ability to promote degradation of nascent DNA at stalled replication forks is required to prevent induction of type I interferons, thereby preventing chronic inflammation (By similarity). Ability to promote DNA end resection at stalled replication forks is independent of dNTPase activity (By similarity). Enhances immunoglobulin hypermutation in B-lymphocytes by promoting transversion mutation (PubMed:29669924).[UniProtKB/Swiss-Prot Function]

**Performance Guaranteed:**

OriGene guarantees that at least two of the three Dicer-Substrate duplexes in the kit will provide at least 70% or more knockdown of the target mRNA when used at 10 nM concentration by quantitative RT-PCR when the TYE-563 fluorescent transfection control duplex (cat# SR30002) indicates that >90% of the cells have been transfected and the HPRT positive control (cat# SR30003) provides 90% knockdown efficiency.

For non-conforming siRNA, requests for replacement product must be made within ninety (90) days from the date of delivery of the siRNA kit. To arrange for a free replacement with newly designed duplexes, please contact Technical Services at [techsupport@origene.com](mailto:techsupport@origene.com). Please provide your data indicating the transfection efficiency and measurement of gene expression knockdown compared to the scrambled siRNA control (quantitative RT-PCR data required).