

Product datasheet for **SR418391**

Ccdc22 Mouse siRNA Oligo Duplex (Locus ID 54638)

Product data:

Product Type:	siRNA Oligo Duplexes
Purity:	HPLC purified
Quality Control:	Tested by ESI-MS
Sequences:	Available with shipment
Stability:	One year from date of shipment when stored at -20°C.
# of transfections:	Approximately 330 transfections/2nmol in 24-well plate under optimized conditions (final conc. 10 nM).
Note:	Single siRNA duplex (10nmol) can be ordered.
RefSeq:	NM_138603
UniProt ID:	Q9JIG7
Synonyms:	AI481216; DXImx40e; Ppp1r3f; Sfc22
Components:	Ccdc22 (Mouse) - 3 unique 27mer siRNA duplexes - 2 nmol each (Locus ID 54638) Included - SR30004, Trilencer-27 Universal Scrambled Negative Control siRNA Duplex - 2 nmol Included - SR30005, RNase free siRNA Duplex Resuspension Buffer - 2 ml
Summary:	Involved in regulation of NF-kappa-B signaling. Promotes ubiquitination of I-kappa-B-kinase subunit IKBKB and its subsequent proteasomal degradation leading to NF-kappa-B activation; the function may involve association with COMMD8 and a CUL1-dependent E3 ubiquitin ligase complex. May down-regulate NF-kappa-B activity via association with COMMD1 and involving a CUL2-dependent E3 ubiquitin ligase complex. Regulates the cellular localization of COMM domain-containing proteins, such as COMMD1 and COMMD10. Component of the CCC complex, which is involved in the regulation of endosomal recycling of surface proteins, including integrins, signaling receptor and channels. The CCC complex associates with SNX17, retriever and WASH complexes to prevent lysosomal degradation and promote cell surface recycling of numerous cargos such as integrins ITGA5:ITGB1. Plays a role in copper ion homeostasis. Involved in copper-dependent ATP7A trafficking between the trans-Golgi network and vesicles in the cell periphery; the function is proposed to depend on its association within the CCC complex and cooperation with the WASH complex on early endosomes.[UniProtKB/Swiss-Prot Function]



[View online »](#)

**Performance
Guaranteed:**

OriGene guarantees that at least two of the three Dicer-Substrate duplexes in the kit will provide at least 70% or more knockdown of the target mRNA when used at 10 nM concentration by quantitative RT-PCR when the TYE-563 fluorescent transfection control duplex (cat# SR30002) indicates that >90% of the cells have been transfected and the HPRT positive control (cat# SR30003) provides 90% knockdown efficiency.

For non-conforming siRNA, requests for replacement product must be made within ninety (90) days from the date of delivery of the siRNA kit. To arrange for a free replacement with newly designed duplexes, please contact Technical Services at techsupport@origene.com. Please provide your data indicating the transfection efficiency and measurement of gene expression knockdown compared to the scrambled siRNA control (quantitative RT-PCR data required).