

Product datasheet for **SR422479**

Kcnma1 Mouse siRNA Oligo Duplex (Locus ID 16531)

Product data:

| | |
|----------------------------|--|
| Product Type: | siRNA Oligo Duplexes |
| Purity: | HPLC purified |
| Quality Control: | Tested by ESI-MS |
| Sequences: | Available with shipment |
| Stability: | One year from date of shipment when stored at -20°C. |
| # of transfections: | Approximately 330 transfections/2nmol in 24-well plate under optimized conditions (final conc. 10 nM). |
| Note: | Single siRNA duplex (10nmol) can be ordered. |
| RefSeq: | NM_001253358 , NM_001253359 , NM_001253360 , NM_001253361 , NM_001253362 , NM_001253363 , NM_001253364 , NM_001253365 , NM_001253366 , NM_001253367 , NM_001253368 , NM_001253369 , NM_001253370 , NM_001253371 , NM_001253372 , NM_001253373 , NM_001253374 , NM_001253375 , NM_001253376 , NM_001253377 , NM_001253378 , NM_010610 |
| UniProt ID: | Q08460 |
| Synonyms: | 5730414M22Rik; BKCa; MaxiK; mSlo; mSlo1; Slo; Slo1 |
| Components: | Kcnma1 (Mouse) - 3 unique 27mer siRNA duplexes - 2 nmol each (Locus ID 16531) Included - SR30004, Trilencer-27 Universal Scrambled Negative Control siRNA Duplex - 2 nmol Included - SR30005, RNase free siRNA Duplex Resuspension Buffer - 2 ml |



[View online »](#)

Summary:

Potassium channel activated by both membrane depolarization or increase in cytosolic Ca(2+) that mediates export of K(+). It is also activated by the concentration of cytosolic Mg(2+). Its activation dampens the excitatory events that elevate the cytosolic Ca(2+) concentration and/or depolarize the cell membrane. It therefore contributes to repolarization of the membrane potential. Plays a key role in controlling excitability in a number of systems, such as regulation of the contraction of smooth muscle, the tuning of hair cells in the cochlea, regulation of transmitter release, and innate immunity. In smooth muscles, its activation by high level of Ca(2+), caused by ryanodine receptors in the sarcoplasmic reticulum, regulates the membrane potential. In cochlea cells, its number and kinetic properties partly determine the characteristic frequency of each hair cell and thereby helps to establish a tonotopic map. Kinetics of KCNMA1 channels are determined by alternative splicing, phosphorylation status and its combination with modulating beta subunits. Highly sensitive to both iberiotoxin (IbTx) and charybdotoxin (CTX).[UniProtKB/Swiss-Prot Function]

Performance Guaranteed:

OriGene guarantees that at least two of the three Dicer-Substrate duplexes in the kit will provide at least 70% or more knockdown of the target mRNA when used at 10 nM concentration by quantitative RT-PCR when the TYE-563 fluorescent transfection control duplex (cat# SR30002) indicates that >90% of the cells have been transfected and the HPRT positive control (cat# SR30003) provides 90% knockdown efficiency.

For non-conforming siRNA, requests for replacement product must be made within ninety (90) days from the date of delivery of the siRNA kit. To arrange for a free replacement with newly designed duplexes, please contact Technical Services at techsupport@origene.com. Please provide your data indicating the transfection efficiency and measurement of gene expression knockdown compared to the scrambled siRNA control (quantitative RT-PCR data required).