

Product datasheet for SR500402

Brcc3 Rat siRNA Oligo Duplex (Locus ID 316794)

Product data:

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product data:	
Product Type:	siRNA Oligo Duplexes
Purity:	HPLC purified
Quality Control:	Tested by ESI-MS
Sequences:	Available with shipment
Stability:	One year from date of shipment when stored at -20°C.
# of transfections:	Approximately 330 transfections/2nmol in 24-well plate under optimized conditions (final conc. 10 nM).
Note:	Single siRNA duplex (10nmol) can be ordered.
RefSeq:	<u>NM 001127300</u>
UniProt ID:	B2RYM5
Components:	Brcc3 (Rat) - 3 unique 27mer siRNA duplexes - 2 nmol each (Locus ID 316794) Included - SR30004, Trilencer-27 Universal Scrambled Negative Control siRNA Duplex - 2 nmol Included - SR30005, RNAse free siRNA Duplex Resuspension Buffer - 2 ml
Summary:	Metalloprotease that specifically cleaves 'Lys-63'-linked polyubiquitin chains. Does not have activity toward 'Lys-48'-linked polyubiquitin chains. Component of the BRCA1-A complex, a complex that specifically recognizes 'Lys-63'-linked ubiquitinated histones H2A and H2AX at DNA lesions sites, leading to target the BRCA1-BARD1 heterodimer to sites of DNA damage at double-strand breaks (DSBs). In the BRCA1-A complex, it specifically removes 'Lys-63'-linked ubiquitin on histones H2A and H2AX, antagonizing the RNF8-dependent ubiquitination at double-strand breaks (DSBs). Catalytic subunit of the BRISC complex, a multiprotein complex that specifically cleaves 'Lys-63'-linked ubiquitin in various substrates. Mediates the specific 'Lys-63'-specific deubiquitination associated with the COP9 signalosome complex (CSN), via the interaction of the BRISC complex with the CSN complex. The BRISC complex is required for normal mitotic spindle assembly and microtubule attachment to kinetochores via its role in deubiquitination of the interferon receptor IFNAR1; deubiquitination increases IFNAR1 activity by enhancing its stability and cell surface expression. Down-regulates the response to bacterial lipopolysaccharide (LPS) via its role in IFNAR1 deubiquitination.[UniProtKB/Swiss-Prot Function]



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2021 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

Performance
Guaranteed:OriGene guarantees that at least two of the three Dicer-Substrate duplexes in the kit will
provide at least 70% or more knockdown of the target mRNA when used at 10 nM
concentration by quantitative RT-PCR when the TYE-563 fluorescent transfection control
duplex (cat# SR30002) indicates that >90% of the cells have been transfected and the HPRT
positive control (cat# SR30003) provides 90% knockdown efficiency.For non-conforming siRNA, requests for replacement product must be made within ninety
(90) days from the date of delivery of the siRNA kit. To arrange for a free replacement with
newly designed duplexes, please contact Technical Services at techsupport@origene.com.
Please provide your data indicating the transfection efficiency and measurement of gene

required).

expression knockdown compared to the scrambled siRNA control (quantitative RT-PCR data

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2021 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US