

## Product datasheet for **SR507553**

### **Mcts1 Rat siRNA Oligo Duplex (Locus ID 302500)**

#### **Product data:**

<b>Product Type:</b>	siRNA Oligo Duplexes
<b>Purity:</b>	HPLC purified
<b>Quality Control:</b>	Tested by ESI-MS
<b>Sequences:</b>	Available with shipment
<b>Stability:</b>	One year from date of shipment when stored at -20°C.
<b># of transfections:</b>	Approximately 330 transfections/2nmol in 24-well plate under optimized conditions (final conc. 10 nM).
<b>Note:</b>	Single siRNA duplex (10nmol) can be ordered.
<b>RefSeq:</b>	<a href="#">NM_001044237</a>
<b>UniProt ID:</b>	<a href="#">Q4G009</a>
<b>Synonyms:</b>	MGC112904
<b>Components:</b>	Mcts1 (Rat) - 3 unique 27mer siRNA duplexes - 2 nmol each (Locus ID 302500) Included - SR30004, Trilencer-27 Universal Scrambled Negative Control siRNA Duplex - 2 nmol Included - SR30005, RNase free siRNA Duplex Resuspension Buffer - 2 ml



[View online »](#)

**Summary:**

Anti-oncogene that plays a role in cell cycle regulation; decreases cell doubling time and anchorage-dependent growth; shortens the duration of G1 transit time and G1/S transition. When constitutively expressed, increases CDK4 and CDK6 kinases activity and CCND1/cyclin D1 protein level, as well as G1 cyclin/CDK complex formation. Plays a role as translation enhancer; Recruits the density-regulated protein/DENR and binds to the cap complex of the 5'-terminus of mRNAs, subsequently altering the mRNA translation profile; Up-regulates protein levels of BCL2L2, TFDP1, MRE11, CCND1 and E2F1, while mRNA levels remains constant. Hyperactivates DNA damage signaling pathway; increased gamma-irradiation-induced phosphorylation of histone H2AX, and induces damage foci formation. Increases the overall number of chromosomal abnormalities such as larger chromosomes formation and multiples chromosomal fusions when overexpressed in gamma-irradiated cells. May play a role in promoting lymphoid tumor development: lymphoid cell lines overexpressing MCTS1 exhibit increased growth rates and display increased protection against apoptosis. May contribute to the pathogenesis and progression of breast cancer via promotion of angiogenesis through the decline of inhibitory THBS1/thrombospondin-1, and inhibition of apoptosis. Involved in the process of proteasome degradation to down-regulate Tumor suppressor p53/TP53 in breast cancer cell; Positively regulates phosphorylation of MAPK1 and MAPK3 (By similarity).[UniProtKB/Swiss-Prot Function]

**Performance Guaranteed:**

OriGene guarantees that at least two of the three Dicer-Substrate duplexes in the kit will provide at least 70% or more knockdown of the target mRNA when used at 10 nM concentration by quantitative RT-PCR when the TYE-563 fluorescent transfection control duplex (cat# SR30002) indicates that >90% of the cells have been transfected and the HPRT positive control (cat# SR30003) provides 90% knockdown efficiency.

For non-conforming siRNA, requests for replacement product must be made within ninety (90) days from the date of delivery of the siRNA kit. To arrange for a free replacement with newly designed duplexes, please contact Technical Services at [techsupport@origene.com](mailto:techsupport@origene.com). Please provide your data indicating the transfection efficiency and measurement of gene expression knockdown compared to the scrambled siRNA control (quantitative RT-PCR data required).