

## **Product datasheet for SR511285**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Morf4l1 Rat siRNA Oligo Duplex (Locus ID 300891)

### **Product data:**

**Product Type:** siRNA Oligo Duplexes

Purity: HPLC purified

Quality Control: Tested by ESI-MS

Sequences: Available with shipment

**Stability:** One year from date of shipment when stored at -20°C.

# of transfections: Approximately 330 transfections/2nmol in 24-well plate under optimized conditions (final

conc. 10 nM).

**Note:** Single siRNA duplex (10nmol) can be ordered.

**RefSeq:** <u>NM 001011999</u>

UniProt ID: Q6AYU1

Components: Morf4l1 (Rat) - 3 unique 27mer siRNA duplexes - 2 nmol each (Locus ID 300891)

Included - SR30004, Trilencer-27 Universal Scrambled Negative Control siRNA Duplex - 2 nmol

Included - SR30005, RNAse free siRNA Duplex Resuspension Buffer - 2 ml

Summary: Component of the NuA4 histone acetyltransferase complex which is involved in

transcriptional activation of select genes principally by acetylation of nucleosomal histones H4 and H2A. This modification may both alter nucleosome - DNA interactions and promote

interaction of the modified histones with other proteins which positively regulate

transcription. This complex may be required for the activation of transcriptional programs

associated with oncogene and proto-oncogene mediated growth induction, tumor

suppressor mediated growth arrest and replicative senescence, apoptosis, and DNA repair. The NuA4 complex ATPase and helicase activities seem to be, at least in part, contributed by the association of RUVBL1 and RUVBL2 with EP400. NuA4 may also play a direct role in DNA repair when directly recruited to sites of DNA damage. Also component of the mSin3A complex which acts to repress transcription by deacetylation of nucleosomal histones. Required for homologous recombination repair (HRR) and resistance to mitomycin C (MMC). Involved in the localization of PALB2, BRCA2 and RAD51, but not BRCA1, to DNA-damage foci

(By similarity).[UniProtKB/Swiss-Prot Function]







# Performance Guaranteed:

OriGene guarantees that at least two of the three Dicer-Substrate duplexes in the kit will provide at least 70% or more knockdown of the target mRNA when used at 10 nM concentration by quantitative RT-PCR when the TYE-563 fluorescent transfection control duplex (cat# SR30002) indicates that >90% of the cells have been transfected and the HPRT positive control (cat# SR30003) provides 90% knockdown efficiency.

For non-conforming siRNA, requests for replacement product must be made within ninety (90) days from the date of delivery of the siRNA kit. To arrange for a free replacement with newly designed duplexes, please contact Technical Services at techsupport@origene.com. Please provide your data indicating the transfection efficiency and measurement of gene expression knockdown compared to the scrambled siRNA control (quantitative RT-PCR data required).