

Product datasheet for SR512156

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Map3k7 Rat siRNA Oligo Duplex (Locus ID 313121)

Product data:

Product Type: siRNA Oligo Duplexes

Purity: HPLC purified

Quality Control: Tested by ESI-MS

Sequences: Available with shipment

Stability: One year from date of shipment when stored at -20°C.

of transfections: Approximately 330 transfections/2nmol in 24-well plate under optimized conditions (final

conc. 10 nM).

Note: Single siRNA duplex (10nmol) can be ordered.

RefSeq: <u>NM 001107920</u>

UniProt ID: P0C8E4
Synonyms: Tak1

Components: Map3k7 (Rat) - 3 unique 27mer siRNA duplexes - 2 nmol each (Locus ID 313121)

Included - SR30004, Trilencer-27 Universal Scrambled Negative Control siRNA Duplex - 2 nmol

Included - SR30005, RNAse free siRNA Duplex Resuspension Buffer - 2 ml

Summary: Serine/threonine kinase which acts as an essential component of the MAP kinase signal

transduction pathway. Plays an important role in the cascades of cellular responses evoked by changes in the environment. Mediates signal transduction of TRAF6, various cytokines including interleukin-1 (IL-1), transforming growth factor-beta (TGFB), TGFB-related factors like BMP2 and BMP4, toll-like receptors (TLR), tumor necrosis factor receptor CD40 and B-cell receptor (BCR). Ceramides are also able to activate MAP3K7/TAK1. Once activated, acts as an upstream activator of the MKK/JNK signal transduction cascade and the p38 MAPK signal transduction cascade through the phosphorylation and activation of several MAP kinase kinases like MAP2K1/MEK1, MAP2K3/MKK3, MAP2K6/MKK6 and MAP2K7/MKK7. These MAP2Ks in turn activate p38 MAPKs, c-jun N-terminal kinases (JNKs) and I-kappa-B kinase complex (IKK). Both p38 MAPK and JNK pathways control the transcription factors activator protein-1 (AP-1), while nuclear factor-kappa B is activated by IKK. MAP3K7 activates also IKBKB and MAPK8/JNK1 in response to TRAF6 signaling and mediates BMP2-induced apoptosis. In osmotic stress signaling, plays a major role in the activation of MAPK8/JNK1, but not that of NF-kappa-B (By similarity). Promotes TRIM5 capsid-specific restriction activity (By similarity).

[UniProtKB/Swiss-Prot Function]







Performance Guaranteed:

OriGene guarantees that at least two of the three Dicer-Substrate duplexes in the kit will provide at least 70% or more knockdown of the target mRNA when used at 10 nM concentration by quantitative RT-PCR when the TYE-563 fluorescent transfection control duplex (cat# SR30002) indicates that >90% of the cells have been transfected and the HPRT positive control (cat# SR30003) provides 90% knockdown efficiency.

For non-conforming siRNA, requests for replacement product must be made within ninety (90) days from the date of delivery of the siRNA kit. To arrange for a free replacement with newly designed duplexes, please contact Technical Services at techsupport@origene.com. Please provide your data indicating the transfection efficiency and measurement of gene expression knockdown compared to the scrambled siRNA control (quantitative RT-PCR data required).