

Product datasheet for SR513590

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Aloxe3 Rat siRNA Oligo Duplex (Locus ID 287424)

Product data:

Product Type: siRNA Oligo Duplexes

Purity: HPLC purified

Quality Control: Tested by ESI-MS

Sequences: Available with shipment

Stability: One year from date of shipment when stored at -20°C.

of transfections: Approximately 330 transfections/2nmol in 24-well plate under optimized conditions (final

conc. 10 nM).

Note: Single siRNA duplex (10nmol) can be ordered.

RefSeq: <u>NM 001105793</u>

UniProt ID: D3ZKX9

Synonyms: e-LOX-3; eLOX-3

Components: Aloxe3 (Rat) - 3 unique 27mer siRNA duplexes - 2 nmol each (Locus ID 287424)

Included - SR30004, Trilencer-27 Universal Scrambled Negative Control siRNA Duplex - 2 nmol

Included - SR30005, RNAse free siRNA Duplex Resuspension Buffer - 2 ml

Summary: Non-heme iron-containing lipoxygenase which is atypical in that it displays a prominent

hydroperoxide isomerase activity and a reduced dioxygenase activity compared to other

lipoxygenases. The hydroperoxide isomerase activity catalyzes the isomerization of

hydroperoxides, derived from arachidonic and linoleic acid by ALOX12B, into hepoxilin-type epoxyalcohols. The dioxygenase activity requires a step of activation of the enzyme by molecular oxygen. In presence of oxygen, oxygenates polyunsaturated fatty acids, including arachidonic acid, to produce fatty acid hydroperoxides. In the skin, acts downstream of ALOX12B on the linoleate moiety of esterified omega-hydroxyacyl-sphingosine (EOS)

ceramides to produce an epoxy-ketone derivative, a crucial step in the conjugation of omegahydroxyceramide to membrane proteins. Therefore plays a crucial role in the synthesis of corneocytes lipid envelope and the establishment of the skin barrier to water loss. In parallel, it may have a signaling function in barrier formation through the production of hepoxilins metabolites. Plays also a role in adipocyte differentiation through hepoxilin A3 and hepoxilin B3 production which in turn activates PPARG. Through the production of hepoxilins in the spinal cord, it may regulate inflammatory tactile allodynia.[UniProtKB/Swiss-Prot Function]





Performance Guaranteed:

OriGene guarantees that at least two of the three Dicer-Substrate duplexes in the kit will provide at least 70% or more knockdown of the target mRNA when used at 10 nM concentration by quantitative RT-PCR when the TYE-563 fluorescent transfection control duplex (cat# SR30002) indicates that >90% of the cells have been transfected and the HPRT positive control (cat# SR30003) provides 90% knockdown efficiency.

For non-conforming siRNA, requests for replacement product must be made within ninety (90) days from the date of delivery of the siRNA kit. To arrange for a free replacement with newly designed duplexes, please contact Technical Services at techsupport@origene.com. Please provide your data indicating the transfection efficiency and measurement of gene expression knockdown compared to the scrambled siRNA control (quantitative RT-PCR data required).