

Product datasheet for SR515092

Acad9 Rat siRNA Oligo Duplex (Locus ID 294973)

Product data:

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product Type:IRNA Oligo DuplexesPurity:H2L purifiedQuality Control:Tested by ESI-MSSequences:Available with shipmentStability:One year from date of shipment when stored at -20°C.# of transfections:Approximately 330 transfections/2nmol in 24-well plate under optimized conditions (final conc. 10 nM).Note:Single siRNA duplex (10nmol) can be ordered.RefSeq:NM 181768UniProt ID:BIVC61Synonyms:Nygg2Components:Acad9 (Rat) - 3 unique 27 mer siRNA duplex es 2 nmol each (Locus ID 294973) Included - SR30004, Trilencer -27 Universal Scrambled Negative Control siRNA Duplex - 2 nmol Included - SR30004, Trilencer -27 Universal Scrambled Negative Control siRNA Duplex - 2 nmol included - SR30004, Trilencer -27 Universal Scrambled Negative Control siRNA Duplex - 2 nmol included - SR30004, Trilencer -27 Universal Scrambled Negative Control siRNA Duplex - 2 nmol included - SR30004, Trilencer -27 Universal Scrambled Negative Control siRNA Duplex - 2 nmol included - SR30004, Trilencer -27 Universal Scrambled Negative Control siRNA Duplex - 2 nmol included - SR30004, Trilencer -27 Universal Scrambled Negative Control siRNA Duplex - 2 nmol included - SR30004, Trilencer -27 Universal Scrambled Negative Control siRNA Duplex - 2 nmol included - SR30004, Trilencer -27 Universal Scrambled Negative Control siRNA Duplex - 2 nmol scrambled Negativ		
Quality Control:Tested by ESI-MSSequences:Available with shipmentStability:One year from date of shipment when stored at -20°C.# of transfections:Approximately 330 transfections/2nmol in 24-well plate under optimized conditions (final conc. 10 nM).Note:Single siRNA duplex (10nmol) can be ordered.RefSeq:NM 181768UniProt ID:B1WC61Synonyms:Nyggf2Components:Acad9 (Rat) - 3 unique 27mer siRNA duplex s - 2 nmol each (Locus ID 294973) Included - SR30004, Trilencer-27 Universal Scrambled Negative Control siRNA Duplex - 2 nmol Included - SR30005, RNAse free siRNA Duplex Resuspension Buffer - 2 mlSummary:As part of the MCIA complex, primarily participates to the assembly of the mitochondrial complex I and therefore plays a role in oxidative phosphorylation. This moonlighting protein has also a dehydrogenase activity toward a broad range of substrates with greater specificity for long-chain unsaturated acyl-CoAs. However, in vivo, it does not seem to play a primary role in fatty acid oxidation. In addition, the function in complex I assembly is independent of	Product Type:	siRNA Oligo Duplexes
Sequences:Available with shipmentStability:One year from date of shipment when stored at -20°C.# of transfections:Approximately 330 transfections/2nmol in 24-well plate under optimized conditions (final conc. 10 nM).Note:Single siRNA duplex (10nmol) can be ordered.RefSeq:NM 181768UniProt ID:B1WC61Synonyms:Nyggf2Components:Acad9 (Rat) - 3 unique 27mer siRNA duplexs - 2 nmol each (Locus ID 294973) Included - SR30004, Trilencer-27 Universal Scrambled Negative Control siRNA Duplex - 2 nmol Included - SR30005, RNAse free siRNA Duplex Resuspension Buffer - 2 mlSummary:As part of the MCIA complex, primarily participates to the assembly of the mitochondrial complex I and therefore plays a role in oxidative phosphorylation. This moonlighting protein has also a dehydrogenase activity toward a broad range of substrates with greater specificity for long-chain unsaturated acyl-CoAs. However, in vivo, it does not seem to play a primary role in fatty acid oxidation. In addition, the function in complex I assembly is independent of	Purity:	HPLC purified
Stability:One year from date of shipment when stored at -20°C.# of transfections:Approximately 330 transfections/2nmol in 24-well plate under optimized conditions (final conc. 10 nM).Note:Single siRNA duplex (10nmol) can be ordered.RefSeq:NM 181768UniProt ID:B1WC61Synonyms:Nyggf2Components:Acad9 (Rat) - 3 unique 27mer siRNA duplexes - 2 nmol each (Locus ID 294973) Included - SR30004, Trilencer-27 Universal Scrambled Negative Control siRNA Duplex - 2 nmol Included - SR30005, RNAse free siRNA Duplex Resuspension Buffer - 2 mlSummary:As part of the MCIA complex, primarily participates to the assembly of the mitochondrial complex I and therefore plays a role in oxidative phosphorylation. This moonlighting protein has also a dehydrogenase activity toward a broad range of substrates with greater specificity for long-chain unsaturated acyl-CoAs. However, in vivo, it does not seem to play a primary role in fatty acid oxidation. In addition, the function in complex I assembly is independent of	Quality Control:	Tested by ESI-MS
# of transfections:Approximately 330 transfections/2nmol in 24-well plate under optimized conditions (final conc. 10 nM).Note:Single siRNA duplex (10nmol) can be ordered.RefSeq:NM 181768UniProt ID:B1WC61Synonyms:Nyggf2Components:Acad9 (Rat) - 3 unique 27mer siRNA duplex s - 2 nmol each (Locus ID 294973) Included - SR30004, Trilencer-27 Universal Scrambled Negative Control siRNA Duplex - 2 nmol Included - SR30005, RNAse free siRNA Duplex Resuspension Buffer - 2 mlSummary:As part of the MCIA complex, primarily participates to the assembly of the mitochondrial complex I and therefore plays a role in oxidative phosphorylation. This moonlighting protein has also a dehydrogenase activity toward a broad range of substrates with greater specificity for long-chain unsaturated acyl-CoAs. However, in vivo, it does not seem to play a primary role in fatty acid oxidation. In addition, the function in complex I assembly is independent of	Sequences:	Available with shipment
conc. 10 nM).Note:Single siRNA duplex (10nmol) can be ordered.RefSeq:NM 181768UniProt ID:B1WC61Synonyms:Nyggf2Components:Acad9 (Rat) - 3 unique 27mer siRNA duplexes - 2 nmol each (Locus ID 294973) Included - SR30004, Trilencer-27 Universal Scrambled Negative Control siRNA Duplex - 2 nmol Included - SR30005, RNAse free siRNA Duplex Resuspension Buffer - 2 mlSummary:As part of the MCIA complex, primarily participates to the assembly of the mitochondrial complex I and therefore plays a role in oxidative phosphorylation. This moonlighting protein has also a dehydrogenase activity toward a broad range of substrates with greater specificity for long-chain unsaturated acyl-CoAs. However, in vivo, it does not seem to play a primary role in fatty acid oxidation. In addition, the function in complex I assembly is independent of	Stability:	One year from date of shipment when stored at -20°C.
RefSeq:NM 181768UniProt ID:B1WC61Synonyms:Nyggf2Components:Acad9 (Rat) - 3 unique 27mer siRNA duplexes - 2 nmol each (Locus ID 294973) Included - SR30004, Trilencer-27 Universal Scrambled Negative Control siRNA Duplex - 2 nmol Included - SR30005, RNAse free siRNA Duplex Resuspension Buffer - 2 mlSummary:As part of the MCIA complex, primarily participates to the assembly of the mitochondrial complex I and therefore plays a role in oxidative phosphorylation. This moonlighting protein has also a dehydrogenase activity toward a broad range of substrates with greater specificity for long-chain unsaturated acyl-CoAs. However, in vivo, it does not seem to play a primary role in fatty acid oxidation. In addition, the function in complex I assembly is independent of	# of transfections:	
UniProt ID:B1WC61Synonyms:Nyggf2Components:Acad9 (Rat) - 3 unique 27mer siRNA duplexes - 2 nmol each (Locus ID 294973) Included - SR30004, Trilencer-27 Universal Scrambled Negative Control siRNA Duplex - 2 nmol Included - SR30005, RNAse free siRNA Duplex Resuspension Buffer - 2 mlSummary:As part of the MCIA complex, primarily participates to the assembly of the mitochondrial complex I and therefore plays a role in oxidative phosphorylation. This moonlighting protein has also a dehydrogenase activity toward a broad range of substrates with greater specificity for long-chain unsaturated acyl-CoAs. However, in vivo, it does not seem to play a primary role in fatty acid oxidation. In addition, the function in complex I assembly is independent of	Note:	Single siRNA duplex (10nmol) can be ordered.
Synonyms:Nyggf2Components:Acad9 (Rat) - 3 unique 27mer siRNA duplexes - 2 nmol each (Locus ID 294973) Included - SR30004, Trilencer-27 Universal Scrambled Negative Control siRNA Duplex - 2 nmol Included - SR30005, RNAse free siRNA Duplex Resuspension Buffer - 2 mlSummary:As part of the MCIA complex, primarily participates to the assembly of the mitochondrial complex I and therefore plays a role in oxidative phosphorylation. This moonlighting protein has also a dehydrogenase activity toward a broad range of substrates with greater specificity for long-chain unsaturated acyl-CoAs. However, in vivo, it does not seem to play a primary role in fatty acid oxidation. In addition, the function in complex I assembly is independent of	RefSeq:	<u>NM 181768</u>
Components:Acad9 (Rat) - 3 unique 27mer siRNA duplexes - 2 nmol each (Locus ID 294973) Included - SR30004, Trilencer-27 Universal Scrambled Negative Control siRNA Duplex - 2 nmol Included - SR30005, RNAse free siRNA Duplex Resuspension Buffer - 2 mlSummary:As part of the MCIA complex, primarily participates to the assembly of the mitochondrial complex I and therefore plays a role in oxidative phosphorylation. This moonlighting protein has also a dehydrogenase activity toward a broad range of substrates with greater specificity for long-chain unsaturated acyl-CoAs. However, in vivo, it does not seem to play a primary role in fatty acid oxidation. In addition, the function in complex I assembly is independent of	UniProt ID:	<u>B1WC61</u>
Included - SR30004, Trilencer-27 Universal Scrambled Negative Control siRNA Duplex - 2 nmol Included - SR30005, RNAse free siRNA Duplex Resuspension Buffer - 2 mlSummary:As part of the MCIA complex, primarily participates to the assembly of the mitochondrial complex I and therefore plays a role in oxidative phosphorylation. This moonlighting protein has also a dehydrogenase activity toward a broad range of substrates with greater specificity for long-chain unsaturated acyl-CoAs. However, in vivo, it does not seem to play a primary role in fatty acid oxidation. In addition, the function in complex I assembly is independent of	Synonyms:	Nyggf2
complex I and therefore plays a role in oxidative phosphorylation. This moonlighting protein has also a dehydrogenase activity toward a broad range of substrates with greater specificity for long-chain unsaturated acyl-CoAs. However, in vivo, it does not seem to play a primary role in fatty acid oxidation. In addition, the function in complex I assembly is independent of	Components:	Included - SR30004, Trilencer-27 Universal Scrambled Negative Control siRNA Duplex - 2 nmol
	Summary:	complex I and therefore plays a role in oxidative phosphorylation. This moonlighting protein has also a dehydrogenase activity toward a broad range of substrates with greater specificity for long-chain unsaturated acyl-CoAs. However, in vivo, it does not seem to play a primary role in fatty acid oxidation. In addition, the function in complex I assembly is independent of



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2021 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	Acad9 Rat siRNA Oligo Duplex (Locus ID 294973) – SR515092
Performance Guaranteed:	OriGene guarantees that at least two of the three Dicer-Substrate duplexes in the kit will provide at least 70% or more knockdown of the target mRNA when used at 10 nM concentration by quantitative RT-PCR when the TYE-563 fluorescent transfection control duplex (cat# SR30002) indicates that >90% of the cells have been transfected and the HPRT positive control (cat# SR30003) provides 90% knockdown efficiency.
	For non-conforming siRNA, requests for replacement product must be made within ninety (90) days from the date of delivery of the siRNA kit. To arrange for a free replacement with newly designed duplexes, please contact Technical Services at techsupport@origene.com. Please provide your data indicating the transfection efficiency and measurement of gene expression knockdown compared to the scrambled siRNA control (quantitative RT-PCR data

required).

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2021 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US