



## MOUSE IGG AGAROSE PRE-CLEARING BEADS

<b>Catalog Number</b>	TA120001
<b>Product Name</b>	Mouse IgG agarose pre-clearing beads
<b>Amount</b>	1mL
<b>Conjugate</b>	Agarose beads
<b>Formulation</b>	TBS (pH7.4) containing 50% Glycerol, 0.02% NaN <sub>3</sub>
<b>Storage/Stability</b>	Store at -20°C. Stable for at least 1 year from date of shipment.
<b>Application</b>	IgG-Agarose can be used as an immunoabsorbent. Pre-clearing the lysate can help reduce non-specific binding of proteins to agarose beads. Pre-clearing with mouse IgG agarose pre-clearing beads will remove proteins that bind immunoglobulins and agarose beads non-specifically. This step will give a lower level of background for immunoprecipitation or protein purification. Optimal working conditions should be determined experimentally by the investigator. Suggested starting dilution 50uL agarose beads conjugated with mouse IgG for pre-clearing for 5-10ug target protein.
<b>Safety</b>	This product contains sodium azide. Sodium azide may react with lead and copper plumbing to form explosive metal azides. Upon disposal of material, flush with a large volume of water to prevent azide accumulation.
<b>Note</b>	This product is for laboratory research use only and is not intended for diagnostic use.
<b>Example of Procedure</b>	<ol style="list-style-type: none"><li>1. Prepare mouse IgG agarose pre-clearing beads by mixing well within the bottle.</li><li>2. Pipette out 50uL mouse IgG agarose pre-clearing beads and transfer to 1.5mL Eppendorf tube.</li><li>3. Spin down the agarose beads briefly at 1,500 rpm and remove the supernatant carefully without disturbing the agarose pellet.</li><li>4. Wash the agarose beads by applying 1mL RIPA buffer and mix well.</li><li>5. Spin down the agarose beads briefly and remove the supernatant carefully without disturbing the agarose pellet.</li><li>6. Repeat step 4 &amp; 5 three times to clean the mouse IgG agarose pre-clearing beads.</li><li>7. Mix the mouse IgG agarose pre-clearing beads with protein mixture sample well (cell lysate expressing target protein and/or negative control lysate) and incubate for 1 hour at 4°C. This step is to pre-clean the sample and avoid the non-specific binding at the following steps.</li><li>8. After pre-clearing, spin down the mouse IgG agarose pre-clearing beads-sample mixture for 1 min at 1,500 rpm. Transfer the supernatant carefully to another new 1.5mL Eppendorf tube. The pre-clearing sample now is ready for immunoprecipitation or protein purification procedure.</li></ol>