

# **Product datasheet for TA150039S**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Mouse Monoclonal tGFP Antibody, Clone OTI2H8, Magnetic Beads

### **Product data:**

**Product Type:** Tag Antibodies

**Product Name:** Mouse Monoclonal tGFP Antibody, Clone OTI2H8, Magnetic Beads

Clone Name: OTI2H8 (formerly 2H8)

Applications: IP, Protein Purification

**Recommended Dilution:** IP: See Datasheet

**Reactivity:** tGFP **Host:** Mouse

Clonality: Monoclonal

Immunogen: Anti-tGFP monoclonal antibody was produced by immunizing mice with purified tGFP protein

expressed in HEK293T cell.

**Isotype:** IgG2b

Formulation: PBS (pH7.4) containing 0.1% BSA, 0.02% NaN3 and 50% glycerol with final concentration

5mg/mL beads

Conjugation: Magnetic Beads

Storage: Shipped at -20°C or with ice packs, Upon delivery store at -20°C. Dilute in PBS(pH7.3) if


necessary. Stable for 12 months from date of receipt. Avoid repeated freeze-thaws.

**Stability:** Stable for 1 year at -20°C from date of shipment.

Predicted Protein Size: 26 kDa



# **Product images:**



Immunoprecipitation (IP) was performed with overexpression lysate expressed with tGFP expression vector (Cat# [PS100010]) or control lysate transfected with pCMV-Entry empty vector (Cat# [PS100001]). Briefly, 20uL anti-tGFP (2H8) magnetic beads (Cat# [TA150039]) were incubated with tGFP overexpression lysate or negative control lysate overnight at 4°C, respectively. After washing three times with RIPA buffer, 20uL 4x SDS sample buffer was applied to the lysate-beads mixture. The samples were boiled at 95°C for 10min. The IP sample supernatants were fractionated on SDS-PAGE gel, blotted onto nitrocellulose membrane. The IP efficiency was evaluated by using HRP-conjugated anti-tGFP antibody (Cat# [TA150043]). Lane 1: anti-tGFP (2H8) magnetic beads (20uL) with tGFP expression lysate; Lane 2: anti-tGFP magnetic beads (20uL) with negative control lysate.