

Product datasheet for TA190146

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com

OriGene Technologies, Inc.

techsupport@origene.com
EU: info-de@origene.com
CN: techsupport@origene.cn

BrdU Mouse Monoclonal Antibody [Clone ID: OTI2B12]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI2B12
Applications: IF, IHC

Recommended Dilution: IHC 1:150, IF 1:150

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Bromodeoxyuridine coupled to keyhole limpet hemocyanin (KLH).

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Note: 5-chloro-2'-deoxyuridine (CIDU), 5-bromo-2'-deoxyuridine (BrdU), 5-iodo-2'-deoxyuridine (IdU)


and 5-ethynyl-2'-deoxyuridine (EdU) are nucleoside analogs of thymidine. Cells that treated with these analogs will incorporate the chemicals into the genomic DNA during S-phase. Immunochemical method detection of these analogs is thus used to quantify the cell proliferation, cell cycle status in vitro or in vivo. Since the thymidine analogs can be passed onto the daughter cells, they can also used to trace dividing cell fate in a short period of time over 3 generations. In addition, combination of different analogs and their specific antibodies

can be used to trace cell fate in different time frames.






Product images:




Immunohistochemical staining of paraffinembedded colon tissue from BrdU injected mouse using anti-BrdU mouse monoclonal antibody. (TA190146). Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.



Immunohistochemical staining of paraffinembedded colon tissue from BrdU injected mouse using anti-BrdU mouse monoclonal antibody. (TA190146). Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.





Immunocytochemistry staining of HT-29 cells pulsed with 5-bromo-2'-deoxyuridine (BrdU)using mouse monoclonal anti-BrdU antibody (1:150)