

Product datasheet for TA300078

Dysbindin (DTNBP1) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 1:1000, IHC: 1:10~50

Reactivity: Human, Mouse

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: This Dtnbp1 antibody is generated from rabbits immunized with a KLH conjugated synthetic

peptide between 296-325 amino acids from the C-terminal region of human Dtnbp1.

Formulation: Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.

Concentration: lot specific

Purification: This antibody is purified through a protein A column, followed by peptide affinity purification.

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Gene Name: dystrobrevin binding protein 1

Database Link: NP 115498

Entrez Gene 94245 MouseEntrez Gene 84062 Human

Q96EV8

Background: Dtnp1 may play a role in organelle biogenesis associated with melanosomes, platelet dense

granules, and lysosomes. A similar protein in mouse is a component of a protein complex termed biogenesis of lysosome-related organelles complex 1 (BLOC-1), and binds to alphaand beta-dystrobrevins, which are components of the dystrophin-associated protein complex (DPC). Mutations in this protein are associated with Hermansky-Pudlak syndrome type 7. This

protein may also be associated with schizophrenia.

Synonyms: BLOC1S8; DBND; HPS7; My031; SDY

Protein Families: Druggable Genome




OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Product images:



Western blot analysis of Dtnbp1 (arrow) using rabbit polyclonal Dysbindin (Dtnbp1) Antibody (Cterm) (Cat.#TA300078).



Formalin-fixed and paraffin-embedded human hepatocarcinoma tissue reacted with Dtnbp1 Antibody (C-term) (Cat.#TA300078), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.