

Product datasheet for TA300151

Product data:

Product Type: Primary Antibodies

REA Rabbit Polyclonal Antibody

Applications: IHC, WB

Recommended Dilution: WB: 1:1000, IHC: 1:10~50

Reactivity: Human (Predicted: Mouse, Rat, Bovine, Chicken)

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: This REA (PHB2) antibody is generated from rabbits immunized with a KLH conjugated

synthetic peptide between 66-95 amino acids from the N-terminal region of human REA

(PHB2).

Formulation: Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.

Concentration: lot specific

Purification: This antibody is purified through a protein G column, followed by dialysis against PBS.

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Database Link: NP 009204

Entrez Gene 12034 MouseEntrez Gene 114766 Rat

Background: PHB2 acts as a mediator of transcriptional repression by nuclear hormone receptors via

recruitment of histone deacetylases. This protein functions as an estrogen receptor (ER)-selective coregulator that potentiates the inhibitory activities of antiestrogens and represses the activity of estrogens. It competes with NCOA1 for modulation of ER transcriptional activity. It is probably involved in regulating mitochondrial respiration activity and in aging.

Synonyms: BAP; Bap37; BCAP37; p22; PNAS-141; REA

Protein Families: Druggable Genome, Transcription Factors




OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Product images:



Western blot analysis of PHB2 (arrow) using rabbit polyclonal PHB2 Antibody (Human C-term) (Cat.#TA300151).



Formalin-fixed and paraffin-embedded human colon carcinoma tissue reacted with PHB2 Antibody (N-term) (Cat.#TA300151), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.