

## **Product datasheet for TA302014S**

## Product datasineet for TAS020143

## **CDK1 Rabbit Polyclonal Antibody**

**Product data:** 

**Product Type:** Primary Antibodies

Applications: WB

Recommended Dilution: WB: 1:1000

**Reactivity:** Human (Predicted: Mouse, Rat, Xenopus)

**Host:** Rabbit

**Isotype:** lg

Clonality: Polyclonal

**Immunogen:** This CDC2 antibody is generated from rabbits immunized with a KLH conjugated synthetic

peptide between 213-241 amino acids from the C-terminal region of human CDC2.

**Formulation:** Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.

**Concentration:** lot specific

**Purification:** This antibody is purified through a protein A column, followed by peptide affinity purification.

**Conjugation:** Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

Predicted Protein Size: 34095 Da

**Gene Name:** cyclin-dependent kinase 1

Database Link: NP 001777

Entrez Gene 12534 MouseEntrez Gene 54237 RatEntrez Gene 983 Human

P06493

**Background:** CDC2 is a member of the Ser/Thr protein kinase family. This protein is a catalytic subunit of

the highly conserved protein kinase complex known as M-phase promoting factor (MPF), which is essential for G1/S and G2/M phase transitions of eukaryotic cell cycle. Mitotic cyclins stably associate with this protein and function as regulatory subunits. The kinase activity of CDC2 is controlled by cyclin accumulation and destruction through the cell cycle. The

phosphorylation and dephosphorylation of this protein also play important regulatory roles

in cell cycle control.

Synonyms: CDC2; CDC28A; P34CDC2



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




**Protein Families:** Druggable Genome, Protein Kinase, Stem cell - Pluripotency

**Protein Pathways:** Cell cycle, Gap junction, Oocyte meiosis, p53 signaling pathway, Progesterone-mediated

oocyte maturation

## **Product images:**



Western blot analysis of CDC2 (arrow) using rabbit polyclonal CDC2 Antibody (C-term) (Cat.# [TA302014]).293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or transiently transfected with the CDC2 gene (Lane 2) (Origene Technologies).