

## **Product datasheet for TA302022**

## **NNMT Rabbit Polyclonal Antibody**

## **Product data:**

**Product Type:** Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 1:1000, IHC: 1:10~50

Reactivity: Human Host: Rabbit

**Isotype:** lg

Clonality: Polyclonal

Immunogen: This NNMT antibody is generated from rabbits immunized with a KLH conjugated synthetic

peptide between 77-106 amino acids from the Central region of human NNMT.

**Formulation:** Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.

**Concentration:** lot specific

**Purification:** This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by

dialysis against PBS.

**Conjugation:** Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 29443 Da

**Gene Name:** nicotinamide N-methyltransferase

Database Link: NP 006160

Entrez Gene 4837 Human

P40261

**Background:** N-methylation is one method by which drug and other xenobiotic compounds are

metabolized by the liver. NNMT is the protein responsible for this enzymatic activity, which

uses S-adenosyl methionine as the methyl donor.

**Synonyms:** nicotinamide N-methyltransferase

**Protein Pathways:** Metabolic pathways, Nicotinate and nicotinamide metabolism




**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



## **Product images:**



Western blot analysis of NNMT (arrow) using rabbit polyclonal NNMT Antibody (Center) (Cat# TA302022). 293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or transiently transfected with the NNMT gene (Lane 2) (Origene Technologies).



Formalin-fixed and paraffin-embedded human hepatocarcinoma tissue reacted with NNMT antibody (Center) (Cat# TA302022), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.