

Product datasheet for TA302033S

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Sigma1 receptor (SIGMAR1) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: FC, WB

Recommended Dilution: WB: 1:1000, FC: 1:10~50

Reactivity: Human (Predicted: Mouse, Rat)

Host: Rabbit

Isotype: lg

Clonality: Polyclonal

Immunogen: This OPRS1 antibody is generated from rabbits immunized with a KLH conjugated synthetic

peptide between 47-81 amino acids from the N-terminal region of human OPRS1.

Formulation: Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.

Purification: This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by

dialysis against PBS.

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 25128 Da

Gene Name: sigma non-opioid intracellular receptor 1

Database Link: NP 005857

Entrez Gene 18391 MouseEntrez Gene 29336 RatEntrez Gene 10280 Human

Q99720

Background: OPRS1 is a receptor protein that interacts with a variety of psychotomimetic drugs, including

cocaine and amphetamines. The receptor is believed to play an important role in the cellular functions of various tissues associated with the endocrine, immune, and nervous systems.


Synonyms: ALS16; DSMA2; hSigmaR1; OPRS1; SIG-1R; sigma1R; SR-BP; SR-BP1; SRBP

Protein Families: Druggable Genome, GPCR, Transmembrane






Product images:



Western blot analysis of OPRS1 (arrow) using rabbit polyclonal OPRS1 Antibody (N-term) (Cat.# [TA302033]).293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or transiently transfected with the OPRS1 gene (Lane 2) (Origene Technologies).



Flow cytometric analysis of NCI-H292 cells using OPRS1 Antibody (N-term) (bottom histogram) compared to a negative control cell (top histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.