

Product datasheet for **TA302104**

RACK1 Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	IHC, WB
Recommended Dilution:	WB: 1:1000, IHC: 1:50~100
Reactivity:	Human (Predicted: Mouse, Rat, Zebrafish, Bovine, Chicken, Pig, Monkey)
Host:	Rabbit
Isotype:	Ig
Clonality:	Polyclonal
Immunogen:	This GNB2L1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 29-58 amino acids from the N-terminal region of human GNB2L1.
Formulation:	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	35076 Da
Gene Name:	receptor for activated C kinase 1
Database Link:	NP_006089 Entrez Gene 14694 Mouse Entrez Gene 83427 Rat Entrez Gene 708526 Monkey Entrez Gene 10399 Human P63244



[View online »](#)

Background:


Protein kinases are enzymes that transfer a phosphate group from a phosphate donor, generally the γ phosphate of ATP, onto an acceptor amino acid in a substrate protein. By this basic mechanism, protein kinases mediate most of the signal transduction in eukaryotic cells, regulating cellular metabolism, transcription, cell cycle progression, cytoskeletal rearrangement and cell movement, apoptosis, and differentiation. With more than 500 gene products, the protein kinase family is one of the largest families of proteins in eukaryotes. The family has been classified in 8 major groups based on sequence comparison of their tyrosine (PTK) or serine/threonine (STK) kinase catalytic domains. The STE group (homologs of yeast Sterile 7, 11, 20 kinases) consists of 50 kinases related to the mitogen-activated protein kinase (MAPK) cascade families (Ste7/MAP2K, Ste11/MAP3K, and Ste20/MAP4K). MAP kinase cascades, consisting of a MAPK and one or more upstream regulatory kinases (MAPKKs) have been best characterized in the yeast pheromone response pathway. Pheromones bind to Ste cell surface receptors and activate yeast MAPK pathway.

Synonyms:


Gnb2-rs1; GNB2L1; H12.3; HLC-7; PIG21

Protein Families:


Druggable Genome

Product images:

The anti-GNB2L1 Pab (Cat. #TA302104) is used in Western blot to detect GNB2L1 in Ramos cell lysate.



Western blot analysis of GNB2L1 (arrow) using GNB2L1 Antibody (N-term) (Cat.#TA302104). 293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or transiently transfected with the GNB2L1 gene (Lane 2) (Origene Technologies).



Formalin-fixed and paraffin-embedded human cancer tissue reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by AEC staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated. BC = breast carcinoma; HC = hepatocarcinoma.