

Product datasheet for TA302184S

Paxillin (PXN) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 1:1000, IHC: 1:10~50

Reactivity: Human Rabbit Host:

Isotype: lg

Clonality: Polyclonal

Immunogen: This PXN antibody is generated from rabbits immunized with a KLH conjugated synthetic

peptide between 94-125 amino acids from human PXN.

Formulation: Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.

Concentration: lot specific

Purification: This antibody is purified through a protein A column, followed by peptide affinity purification.

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 64533 Da Gene Name: paxillin Database Link:

NP 002850

Entrez Gene 5829 Human

P49023

Background: PXN is a cytoskeletal adapter protein involved in organisation and function of focal

> adhesions, which are critical to cell adhesion and migration. This in turn plays a role in a wide variety of processes including embryogenesis, organogenesis, wound repair, inflammation and cancer. PXN contains LD motifs, LIM domains, SH3 and SH2 binding domains that serve

as docking sites for cytoskeletal proteins, tyrosine kinases (e.g., FAK, Pyk 2, Src),

serine/threonine kinases, GTPase activating proteins and other adaptor proteins (e.g., Actin,

Vinculin, Crk).

Synonyms: FLJ16691



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Protein Families: Druggable Genome, Stem cell - Pluripotency


Protein Pathways: Chemokine signaling pathway, Focal adhesion, Leukocyte transendothelial migration,

Regulation of actin cytoskeleton, VEGF signaling pathway

Product images:



Western blot analysis of PXN (arrow) using rabbit polyclonal PXN Antibody (Y118) (Cat.# [TA302184]). 293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or transiently transfected with PXN gene (Lane 2) (Origene Technologies).



Formalin-fixed and paraffin-embedded human prostata carcinoma tissue reacted with PXN Antibody (Y118) (Cat.#[TA302184]), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.