

Product datasheet for **TA302224S**

TRPM8 Rabbit Polyclonal Antibody

Product data:


Product Type:	Primary Antibodies
Applications:	IHC, WB
Recommended Dilution:	WB: 1:1000, IHC: 1:10~50
Reactivity:	Human (Predicted: Mouse)
Host:	Rabbit
Isotype:	Ig
Clonality:	Polyclonal
Immunogen:	This TRPM8 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 926-956 amino acids from the C-terminal region of human TRPM8.
Formulation:	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.
Concentration:	lot specific
Purification:	This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	127685 Da
Gene Name:	transient receptor potential cation channel subfamily M member 8
Database Link:	NP_076985 Entrez Gene 171382 Mouse Entrez Gene 79054 Human Q7Z2W7
Background:	TRPM8 is a receptor-activated non-selective cation channel involved in detection of sensations such as coolness, by being activated by cold temperature below 25 degrees Celsius. It is activated by icilin, eucalyptol, menthol, cold and modulation of intracellular pH and involved in menthol sensation. It is permeable for monovalent cations sodium, potassium, and cesium and divalent cation calcium.
Synonyms:	LTRPC6; TRPP8




[View online »](#)

Protein Families: Druggable Genome, Ion Channels: Transient receptor potential, Transmembrane

Product images:



Western blot analysis of TRPM8 (arrow) using rabbit polyclonal TRPM8 Antibody (C-term C940) (Cat.#[TA302224]). 293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or transiently transfected with the TRPM8 gene (Lane 2) (Origene Technologies).



Formalin-fixed and paraffin-embedded human prostate carcinoma tissue reacted with TRPM8 antibody (C-term C940) (Cat.#[TA302224]), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.