

# **Product datasheet for TA302463**

# **DISC1 Goat Polyclonal Antibody**

**Product data:** 

**Product Type:** Primary Antibodies

Applications: WB

**Recommended Dilution:** ELISA: 1:128,000. WB: 0.1-0.3µg/ml.

Reactivity: Human
Host: Goat
Isotype: IgG

Clonality: Polyclonal

**Immunogen:** Peptide with sequence C-HSEDKRKTPLKVLE, from the internal region of the protein sequence

according to NP\_001012975.1; NP\_061132.2.

**Formulation:** Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum

albumin.

**Concentration:** lot specific

**Purification:** Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity

chromatography using the immunizing peptide. Supplied at 0.5 mg/ml in Tris saline, 0.02%

sodium azide, pH7.3 with 0.5% bovine serum albumin.

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

Predicted Protein Size: 97244 Da

**Gene Name:** disrupted in schizophrenia 1

Database Link: NP 061132

Entrez Gene 27185 Human

Q9NRI5



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com

### **DISC1 Goat Polyclonal Antibody - TA302463**

Background:

This gene encodes a protein with multiple coiled coil motifs which is located in the nucleus, cytoplasm and mitochondria. The protein is involved in neurite outgrowth and cortical development through its interaction with other proteins. This gene is disrupted in a t(1;11) (q42.1;q14.3) translocation which segregates with schizophrenia and related psychiatric disorders in a large Scottish family. Alternate transcriptional splice variants, encoding different isoforms, have been characterized. [provided by RefSeq]

Synonyms: C1orf136; SCZD9

## **Product images:**



TA302463 (0.1ug/ml) staining of human brain (hippocampus) lysate (35ug protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.