

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Product datasheet for TA302496

# **GCNT3 Goat Polyclonal Antibody**

### **Product data:**

Product Type:	Primary Antibodies
Applications:	WB
Recommended Dilution:	ELISA: 1:2,000. WB: 0.3-1µg/ml.
Reactivity:	Human
Host:	Goat
lsotype:	lgG
Clonality:	Polyclonal
Immunogen:	Peptide with sequence C-RDTLHLTNKKKD, from the internal region of the protein sequence according to NP_004742.1.
Formulation:	Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.
Concentration:	lot specific
Purification:	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide. Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	54490 Da
Gene Name:	glucosaminyl (N-acetyl) transferase 3, mucin type
Database Link:	<u>NP_004742</u> <u>Entrez Gene 9245 Human</u> <u>O95395</u>
Background:	This enzyme catalyzes O-glycan branch synthesis of the core 2 and core 4 type in mucins and controls expression of core 2 branched oligosaccharides and I antigens on the cell surface. [provided by RefSeq]. COMPLETENESS: complete on the 3' end.
Synonyms:	4GnT; C2; C2GNT2; C2GNTM; C24GNT; GNTM



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

# **GCNT3** Goat Polyclonal Antibody – TA302496

Protein Families: Transmembrane

Protein Pathways: Metabolic pathways, O-Glycan biosynthesis

# **Product images:**

 250kDa 150kDa 100kDa 75kDa 50kDa
37kDa 25kDa
20kDa

15kDa

TA302496 (0.3ug/ml) staining of human lymph node lysate (35ug protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US