

Product datasheet for TA302687

OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

RANGAP1 Goat Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: WB

Recommended Dilution: ELISA: 1:32,000. WB 0.1-0.5μg/ml.

Reactivity: Human, Mouse (Expected from sequence similarity: Rat, Dog, Cow)

Host: Goat Isotype: IgG

Clonality: Polyclonal

Immunogen: Peptide with sequence ASEDIAKLAETLAK-C, from the N Terminus of the protein sequence

according to NP_002874.

Formulation: Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum

albumin.

Concentration: lot specific

Purification: Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity

chromatography using the immunizing peptide. Supplied at 0.5 mg/ml in Tris saline, 0.02%

sodium azide, pH7.3 with 0.5% bovine serum albumin.

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 63.4 kDa

Gene Name: Ran GTPase activating protein 1

Database Link: NP 002874

Entrez Gene 19387 MouseEntrez Gene 362965 RatEntrez Gene 481234 DogEntrez Gene 5905

<u>Human</u> P46060






Background:

RanGAP1, is a homodimeric 65-kD polypeptide that specifically induces the GTPase activity of RAN, but not of RAS by over 1,000-fold. RanGAP1 is the immediate antagonist of RCC1, a regulator molecule that keeps RAN in the active, GTP-bound state. The RANGAP1 gene encodes a 587-amino acid polypeptide. The sequence is unrelated to that of GTPase activators for other RAS-related proteins, but is 88% identical to Fug1, the murine homolog of yeast Rna1p. RanGAP1 and RCC1 control RAN-dependent transport between the nucleus and cytoplasm. RanGAP1 is a key regulator of the RAN GTP/GDP cycle. [provided by RefSeq]

Synonyms: Fug1; RANGAP; SD

Product images:



TA302687 staining (0.2ug/ml) of 3T3 lysate (RIPA buffer, 35ug total protein per lane). Primary incubated for 1 hour. Detected by western blot using chemiluminescence.