

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Product datasheet for TA302754

## **DDAH1 Goat Polyclonal Antibody**

### **Product data:**

| Product Type:         | Primary Antibodies   |
|-----------------------|--|
| Applications:         | WB   |
| Recommended Dilution: | ELISA: 1:2,000. WB: 1-3µg/ml.  |
| Reactivity:           | Human (Expected from sequence similarity: Cow)   |
| Host:                 | Goat   |
| lsotype:              | lgG  |
| Clonality:            | Polyclonal   |
| Immunogen:            | Peptide with sequence TCCSVLINKKVDS, from the C Terminus of the protein sequence according to NP_036269.1.   |
| Formulation:          | Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.  |
| Concentration:        | lot specific   |
| Purification:         | Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide. Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing. |
| Conjugation:          | Unconjugated   |
| Storage:              | Store at -20°C as received.  |
| Stability:            | Stable for 12 months from date of receipt.   |
| Gene Name:            | dimethylarginine dimethylaminohydrolase 1  |
| Database Link:        | <u>NP_001127917</u><br><u>Entrez Gene 23576 Human</u><br><u>O94760</u>   |
| Background:           | This gene belongs to the dimethylarginine dimethylaminohydrolase (DDAH) gene family. The encoded enzyme plays a role in nitric oxide generation by regulating cellular concentrations of methylarginines, which in turn inhibit nitric oxide synthase activity. [provided by RefSeq]             |
| Synonyms:             | DDAH; DDAH-1; DDAHI; HEL-S-16  |



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

# **Product images:**

| 250kDa<br>150kDa<br>100kDa<br>75kDa |  |
|-------------------------------------|--|
| 50kDa                               |  |
| 🗩 37kDa                             | TA302754 staining (0.1ug/ml) of human kidney<br>lysate (RIPA buffer, 35ug total protein per lane).<br>Primary incubated for 1 hour. Detected by<br>western blot using chemiluminescence. |
| 25kDa<br>20kDa                      |  |
| 15kDa                               |  |
| 10kDa                               |  |

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US