

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Product datasheet for TA302864

## **RNF87 (TRIM4) Goat Polyclonal Antibody**

## **Product data:**

Product Type:	Primary Antibodies	
Applications:	WB	
Recommended Dilution:	ELISA: 1:32,000. WB: 1µg/ml.	
Reactivity:	Human	
Host:	Goat	
lsotype:	lgG	
Clonality:	Polyclonal	
Immunogen:	Peptide with sequence C-ASLVIPPVTDRK, from the C Terminus of the protein sequence according to NP_148977; NP_149082.	
Formulation:	Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.	
Concentration:	lot specific	
Purification:	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide. Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing.	
Conjugation:	Unconjugated	
Storage:	Store at -20°C as received.	
Stability:	Stable for 12 months from date of receipt.	
Gene Name:	tripartite motif containing 4	
Database Link:	<u>NP_148977</u> <u>Entrez Gene 89122 Human</u> <u>Q9C037</u>	
Background:	The protein encoded by this gene is a member of the tripartite motif (TRIM) family. The TRIM motif includes three zinc-binding domains, a RING, a B-box type 1 and a B-box type 2, and a coiled-coil region. The protein localizes to cytoplasmic bodies. Its function has not been identified. Alternative splicing of this gene generates two transcript variants. [provided by RefSeq]	



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	87 (TRIM4) Goat Polyclonal Antibody – T	A302864
Synonyms:	RNF87	
Protein Families:	Druggable Genome	
Product images:		
	250kDa 150kDa 100kDa 75kDa	TA302864 staining (1ug/ml) of HepG2 lysate (RIPA
	50kDa 37kDa	buffer, 30ug total protein per lane). Primary incubated for 1 hour. Detected by western blot using chemiluminescence.
	25kDa	

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US