

## Product datasheet for **TA302886**


### ETEA (FAF2) Goat Polyclonal Antibody

#### Product data:

Product Type:	Primary Antibodies
Applications:	WB
Recommended Dilution:	ELISA: 1:4,000. WB: 1-3µg/ml.
Reactivity:	Human (Expected from sequence similarity: Mouse, Rat, Dog, Cow)
Host:	Goat
Isotype:	IgG
Clonality:	Polyclonal
Immunogen:	Peptide with sequence C-HTEVLFVQDLTDE, from the C Terminus of the protein sequence according to NP_055428.1.
Formulation:	Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.
Purification:	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide. Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing.
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Gene Name:	Fas associated factor family member 2
Database Link:	<a href="#">NP_055428</a> <a href="#">Entrez Gene 76577 Mouse</a> <a href="#">Entrez Gene 291000 Rat</a> <a href="#">Entrez Gene 489102 Dog</a> <a href="#">Entrez Gene 23197 Human</a> <a href="#">Q96CS3</a>
Background:	The protein encoded by this gene is highly expressed in peripheral blood of patients with atopic dermatitis (AD), compared to normal individuals. It may play a role in regulating the resistance to apoptosis that is observed in T cells and eosinophils of AD patients. [provided by RefSeq]. COMPLETENESS: complete on the 3' end.
Synonyms:	ETEa; UBxD8; UBxN3B



[View online »](#)

**Product images:**

TA302886 staining (1ug/ml) of Human Testis lysate (RIPA buffer, 35ug total protein per lane). Primary incubated for 1 hour. Detected by western blot using chemiluminescence.