

## Product datasheet for **TA302917**

### KPNA2 Goat Polyclonal Antibody

#### Product data:

Product Type:	Primary Antibodies
Applications:	IF, WB
Recommended Dilution:	WB: 1-3 µg/ml
Reactivity:	Human (Expected from sequence similarity: Mouse, Rat, Dog)
Host:	Goat
Isotype:	IgG
Clonality:	Polyclonal
Immunogen:	Peptide with sequence C-QVQDGAPGTFNF, from the C Terminus of the protein sequence according to NP_002257.1.
Formulation:	Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.
Concentration:	lot specific
Purification:	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide. Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing.
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Gene Name:	karyopherin subunit alpha 2
Database Link:	<a href="#">NP_002257</a> <a href="#">Entrez Gene 16647 Mouse</a> <a href="#">Entrez Gene 85245 Rat</a> <a href="#">Entrez Gene 480469 Dog</a> <a href="#">Entrez Gene 3838 Human</a> <a href="#">P52292</a>



[View online »](#)

**Background:**


The import of proteins into the nucleus is a process that involves at least 2 steps. The first is an energy-independent docking of the protein to the nuclear envelope and the second is an energy-dependent translocation through the nuclear pore complex. Imported proteins require a nuclear localization sequence (NLS) which generally consists of a short region of basic amino acids or 2 such regions spaced about 10 amino acids apart. Proteins involved in the first step of nuclear import have been identified in different systems. These include the *Xenopus* protein importin and its yeast homolog, SRP1 (a suppressor of certain temperature-sensitive mutations of RNA polymerase I in *Saccharomyces cerevisiae*), which bind to the NLS. KPNA2 protein interacts with the NLSs of DNA helicase Q1 and SV40 T antigen and may be involved in the nuclear transport of proteins. KPNA2 also may play a role in V(D)J recombination [provided by RefSeq].

**Synonyms:**


IPOA1; QIP2; RCH1; SRP1-alpha; SRP1alpha

**Protein Families:**

Druggable Genome, Stem cell - Pluripotency

**Product images:**

TA302917 staining (0.1 ug/ml) of HeLa lysate (RIPA buffer, 35 ug total protein per lane). Primary incubated for 1 hour. Detected by western blot using chemiluminescence.



TA302917 staining (top panel and green) of 4% formaldehyde-fixed HeLa with endogenously expressed HistonH2B-GFP fusion as DNA marker (mid panel and red).