

Product datasheet for TA303061

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Cannabinoid Receptor II (CNR2) Goat Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: WB

Recommended Dilution: ELISA: 1:16,000. WB: 2-3µg/ml.

Reactivity: Human
Host: Goat
Isotype: IgG

Clonality: Polyclonal

Immunogen: Peptide with sequence C-TETEADGKITPWPD, from the C Terminus of the protein sequence

according to NP_001832.1.

Formulation: Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum

albumin.

Concentration: lot specific

Purification: Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity

chromatography using the immunizing peptide. Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Aliquot and store at -20 $^{\circ}$ C. Minimize

freezing and thawing.

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Gene Name: cannabinoid receptor 2

Database Link: NP 001832

Entrez Gene 1269 Human

P34972





Cannabinoid Receptor II (CNR2) Goat Polyclonal Antibody - TA303061

the G-protein-coupled receptors. [provided by RefSeq]

Background: The cannabinoid delta-9-tetrahydrocannabinol is the principal psychoactive ingredient of

marijuana. The proteins encoded by this gene and the cannabinoid receptor 1 (brain) (CNR1) gene have the characteristics of a guanine nucleotide-binding protein (G-protein)-coupled receptor for cannabinoids. They inhibit adenylate cyclase activity in a dose-dependent, stereoselective, and pertussis toxin-sensitive manner. These proteins have been found to be involved in the cannabinoid-induced CNS effects (including alterations in mood and cognition) experienced by users of marijuana. The cannabinoid receptors are members of family 1 of

Synonyms: CB-2; CB2; CX5

Protein Families: Druggable Genome, GPCR, Transmembrane

Protein Pathways: Neuroactive ligand-receptor interaction

Product images:



TA303061 (2ug/ml) staining of Jurkat lysate (35ug protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.