

## **Product datasheet for TA303072**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## **Mybl2 Goat Polyclonal Antibody**

**Product data:** 

**Product Type:** Primary Antibodies

Applications: WB

**Recommended Dilution:** ELISA: 1:32,000. WB: 0.1-0.3µg/ml.

**Reactivity:** Rat (Expected from sequence similarity: Human, Mouse, Dog, Pig, Cow)

Host: Goat Isotype: IgG

Clonality: Polyclonal

**Immunogen:** Peptide with sequence C-QEKARQLLGRLKPSH, from the C Terminus of the protein sequence

according to NP\_002457.1.

**Formulation:** Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum

albumin.

**Concentration:** lot specific

**Purification:** Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity

chromatography using the immunizing peptide. Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize

freezing and thawing.

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Gene Name:** myeloblastosis oncogene-like 2

Database Link: NP 001100006

Entrez Gene 4605 HumanEntrez Gene 17865 MouseEntrez Gene 477232 DogEntrez Gene

296344 Rat





Background: The protein encoded by this gene, a member of the MYB family of transcription factor genes,

is a nuclear protein involved in cell cycle progression. The encoded protein is phosphorylated by cyclin A/cyclin-dependent kinase 2 during the S-phase of the cell cycle and possesses both activator and repressor activities. It has been shown to activate the cell division cycle 2, cyclin D1, and insulin-like growth factor-binding protein 5 genes. Transcript variants may exist for this gene, but their full-length natures have not been determined. [provided by RefSeq]

B-MYB; BMYB; MGC15600 Synonyms:

## **Product images:**



TA303072 (0.1ug/ml) staining of rat stomach lysate (35ug protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.