

Product datasheet for TA303081

PARP2 Goat Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: WB

Recommended Dilution: ELISA: 1:8,000. WB: 1-3µg/ml.

Reactivity: Human
Host: Goat
Isotype: IgG

Clonality: Polyclonal

Immunogen: Peptide with sequence C-LDLFEVEKDGEKE, from the internal region of the protein sequence

according to NP_005475.1.

Formulation: Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum

albumin.

Purification: Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity

chromatography using the immunizing peptide. Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize

freezing and thawing.

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Gene Name: poly(ADP-ribose) polymerase 2

Database Link: NP 001036083

Entrez Gene 10038 Human

Q9UGN5



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com

PARP2 Goat Polyclonal Antibody - TA303081


Background: This gene encodes poly(ADP-ribosyl)transferase-like 2 protein, which contains a catalytic

domain and is capable of catalyzing a poly(ADP-ribosyl)ation reaction. This protein has a catalytic domain which is homologous to that of poly (ADP-ribosyl) transferase, but lacks an N-terminal DNA binding domain which activates the C-terminal catalytic domain of poly (ADP-ribosyl) transferase. The basic residues within the N-terminal region of this protein may bear potential DNA-binding properties, and may be involved in the nuclear and/or nucleolar targeting of the protein. Two alternatively spliced transcript variants encoding distinct isoforms have been found. [provided by RefSeq]

Synonyms: ADPRTL2; ADPRTL3; ARTD2; pADPRT-2; PARP-2

Protein Families: Druggable Genome
Protein Pathways: Base excision repair

Product images:



TA303081 (1ug/ml) staining of Human Spleen Lysate (35ug protein in RIPA buffer). Primary incubation was 1 hour. Detected by

chemiluminescence.