

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Product datasheet for TA303083

## **ARHGEF5 Goat Polyclonal Antibody**

## **Product data:**

Product Type:	Primary Antibodies	
Applications:	WB	
Recommended Dilution:	ELISA: 1:16,000. WB: 0.1-0.3µg/ml.	
Reactivity:	Human	
Host:	Goat	
lsotype:	lgG	
Clonality:	Polyclonal	
Immunogen:	Peptide with sequence C-DKEIDQNSQQEE, from the internal region of the protein sequence according to NP_005426.2.	
Formulation:	Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.	
Concentration:	lot specific	
Purification:	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide. Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing.	
Conjugation:	Unconjugated	
Storage:	Store at -20°C as received.	
Stability:	Stable for 12 months from date of receipt.	
Gene Name:	Rho guanine nucleotide exchange factor 5	
Database Link:	<u>NP_005426</u> <u>Entrez Gene 7984 Human</u> <u>Q12774</u>	
Background:	Rho GTPases play a fundamental role in numerous cellular processes initiated by extracellular stimuli that work through G protein coupled receptors. The encoded protein may form a complex with G proteins and stimulate Rho-dependent signals. This protein may be involved in the control of cytoskeletal organization. [provided by RefSeq]. COMPLETENESS: complete on the 3' end.	



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	ARHGEF5 Goat Polyclonal Antibody – TA303083	
Synonyms:	GEF5; P60; TIM; TIM1	
Protein Families:	Druggable Genome	
Product image	es:	
	250kDa 150kDa 100kDa 75kDa 50kDa 37kDa 25kDa 20kDa 15kDa 10kDa	TA303083 (0.1ug/ml) staining of HeLa Lysate (35ug protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US