

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Product datasheet for TA303153

## **GATA5 Goat Polyclonal Antibody**

## Product data:

Product Type:	Primary Antibodies	
Applications:	WB	
Recommended Dilution:	ELISA: 1:32,000. WB: 0.3-1µg/ml.	
Reactivity:	Human	
Host:	Goat	
lsotype:	IgG	
Clonality:	Polyclonal	
Immunogen:	Peptide with sequence C-ELAARPGWAQTATAD, from the internal region of the protein sequence according to NP_536721.1.	
Formulation:	Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.	
Concentration:	lot specific	
Purification:	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide. Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing.	
Conjugation:	Unconjugated	
Storage:	Store at -20°C as received.	
Stability:	Stable for 12 months from date of receipt.	
Gene Name:	GATA binding protein 5	
Database Link:	<u>NP_536721</u> <u>Entrez Gene 140628 Human</u> <u>Q9BWX5</u>	
Background:	The protein encoded by this gene is a transcription factor that contains two GATA-type zinc fingers. The encoded protein is known to bind to hepatocyte nuclear factor-1alpha (HNF-1alpha), and this interaction is essential for cooperative activation of the intestinal lactase-phlorizin hydrolase promoter. In other organisms, similar proteins may be involved in the establishment of cardiac smooth muscle cell diversity.	



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	GATA5 Goat Polyclonal Antibody – TA303153	
Synonyms:	bB379O24.1; GATAS	
Protein Families:	Transcription Factors	
Product image	s:	
	250kDa 150kDa 100kDa 75kDa 50kDa 37kDa 25kDa 20kDa 15kDa 10kDa	TA303153 (0.3ug/ml) staining of HepG2 cell lysate (35ug protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US