

Product datasheet for TA303162

HOXB13 Goat Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: WB

Recommended Dilution: ELISA: 1:128,000. WB: 0.1-0.3µg/ml.

Reactivity: Human
Host: Goat
Isotype: IgG

Clonality: Polyclonal

Immunogen: Peptide with sequence C-KKVLAKVKNSATP, from the C Terminus of the protein sequence

according to NP_006352.2.

Formulation: Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum

albumin.

Concentration: lot specific

Purification: Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity

chromatography using the immunizing peptide. Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize

freezing and thawing.

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Gene Name: homeobox B13

Database Link: NP 006352

Entrez Gene 10481 Human

Q92826



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



HOXB13 Goat Polyclonal Antibody - TA303162


Background: This gene encodes a transcription factor that belongs to the homeobox gene family. Genes of

this family are highly conserved among vertebrates and essential for vertebrate embryonic development. This gene has been implicated to play a role in fetal skin development and cutaneous regeneration. In mice, a similar gene was shown to exhibit temporal and spatial colinearity in the main body axis of the embryo, but was not expressed in the secondary axes, which suggests functions in body patterning along the axis. This gene and other HOXB genes form a gene cluster at chromosome the 17q21-22 region. [provided by RefSeq]

Synonyms: PSGD

Protein Families: Transcription Factors

Product images:



TA303162 (0.1ug/ml) staining of HeLa cell lysate (35ug protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.