

## Product datasheet for **TA303191**

### FAIM1 (FAIM) Goat Polyclonal Antibody

#### Product data:

Product Type:	Primary Antibodies
Applications:	WB
Recommended Dilution:	ELISA: 1:32,000. WB: 0.5-1 µg/ml.
Reactivity:	Human (Expected from sequence similarity: Mouse, Rat, Dog)
Host:	Goat
Isotype:	IgG
Clonality:	Polyclonal
Immunogen:	Peptide with sequence C-KKYMEDRSKTTN, from the internal region of the protein sequence according to NP_001028202.1; NP_001028203.1; NP_001028204.1; NP_060617.1.
Formulation:	Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.
Concentration:	lot specific
Purification:	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide. Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing.
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Gene Name:	Fas apoptotic inhibitory molecule
Database Link:	<a href="#">NP_001028202</a> <a href="#">Entrez Gene 23873</a> <a href="#">MouseEntrez Gene 140930</a> <a href="#">RatEntrez Gene 485687</a> <a href="#">DogEntrez Gene 55179</a> <a href="#">Human</a> <a href="#">Q9NVQ4</a>
Background:	Plays a role as an inducible effector molecule that mediates Fas resistance produced by surface Ig engagement in B cells (By similarity)
Synonyms:	FAIM1



[View online »](#)

**Product images:**

TA303191 (0.5ug/ml) staining of Human Lymph Node lysate (35ug protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.