

Product datasheet for TA303341

Dach2 Goat Polyclonal Antibody

Product data:

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

| Product Type: | Primary Antibodies |
|-----------------------|--|
| Applications: | WB |
| Recommended Dilution: | ELISA: 1:64,000. WB: 0.03-0.1µg/ml. |
| Reactivity: | Mouse (Expected from sequence similarity: Human) |
| Host: | Goat |
| lsotype: | lgG |
| Clonality: | Polyclonal |
| Immunogen: | Peptide with sequence C-TRKQAVNSSRPGR, from the internal region of the protein sequence according to NP_444511.1. |
| Formulation: | Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. |
| Concentration: | lot specific |
| Purification: | Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide. Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing. |
| Conjugation: | Unconjugated |
| Storage: | Store at -20°C as received. |
| Stability: | Stable for 12 months from date of receipt. |
| Gene Name: | dachshund 2 (Drosophila) |
| Database Link: | <u>NP_291083</u> Entrez Gene 117154 HumanEntrez Gene 93837 Mouse Q925Q8 |



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

Dach2 Goat Polyclonal Antibody – TA303341

Background: This gene is one of two genes which encode a protein similar to the Drosophila protein dachshund, a transcription factor involved in cell fate determination in the eye, limb and genital disc of the fly. The encoded protein contains two characteristic dachshund domains: an N-terminal domain responsible for DNA binding and a C-terminal domain responsible for protein-protein interactions. This gene is located on the X chromosome and is subject to inactivation by DNA methylation. The encoded protein may be involved in regulation of organogenesis and myogenesis, and may play a role in premature ovarian failure. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq]

Synonyms: FLJ31391; MGC138545; OTTHUMP00000023630

25040-

Product images:

| 250kDa 150kDa 100kDa 75kDa | |
|-------------------------------------|---|
| 50kDa | TA303341 (0.03ug/ml) staining of NIH3T3 cell lysate (35ug protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence. |
| 37kDa | |
| 25kDa | |
| 20kDa | |
| 15kDa | |

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US