

# Product datasheet for TA305652

## DRAK2 (STK17B) Goat Polyclonal Antibody

### **Product data:**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product Type:	Primary Antibodies
Applications:	WB
Recommended Dilution:	WB: 1-3ug/ml.
Reactivity:	Human (Expected from sequence similarity: Dog)
Host:	Goat
lsotype:	IgG
Clonality:	Polyclonal
Immunogen:	Peptide with sequence C-HSVRSSEDKTSK, from the internal region of the protein sequence according to NP_004217.1.
Formulation:	0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin
Concentration:	lot specific
Purification:	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide. Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Aliquot and store at -20C. Minimize freezing and thawing.
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Gene Name:	serine/threonine kinase 17b
Database Link:	<u>NP_004217</u> <u>Entrez Gene 488453 DogEntrez Gene 9262 Human</u> <u>O94768</u>
Synonyms:	DRAK2
Protein Families:	Druggable Genome, Protein Kinase



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

## **Product images:**

250kDa 150kDa 100kDa 75kDa	
50kDa	
37kDa	
	TA305652 (1ug/ml) staining of MOLT4 lysate
25kDa	(35ug protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.
20kDa	was i nour. Detected by chemilaninescence.
15kDa	

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US