

Product datasheet for TA305735

TMPRSS4 Goat Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: WB

Recommended Dilution: WB: 0.3-1ug/ml.

Reactivity: Human
Host: Goat
Isotype: IgG

Clonality: Polyclonal

Immunogen: Peptide with sequence HCFRKHTDVFNWK, from the internal region of the protein sequence

according to NP_063947.1; NP_001077416.1.

Formulation: 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

Concentration: lot specific

Purification: Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity

chromatography using the immunizing peptide. Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Aliquot and store at -20C. Minimize

freezing and thawing.

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Gene Name: transmembrane protease, serine 4

Database Link: NP 001077416

Entrez Gene 56649 Human

Q9NRS4

Background: This gene encodes a member of the serine protease family. Serine proteases are known to be

involved in a variety of biological processes, whose malfunction often leads to human diseases and disorders. This gene was identified as a gene overexpressed in pancreatic carcinoma. The encoded protein is membrane bound with a N-terminal anchor sequence and a glycosylated extracellular region containing the serine protease domain. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq]



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




TMPRSS4 Goat Polyclonal Antibody - TA305735

Synonyms: CAPH2; MT-SP2; TMPRSS3

Protein Families: Druggable Genome, Protease, Transmembrane

Product images:



TA305735 (1ug/ml) staining of HEK293 lysate (35ug protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.