

Product datasheet for **TA305768**


RIP140 (NRIP1) Goat Polyclonal Antibody

Product data:

| | |
|------------------------------|---|
| Product Type: | Primary Antibodies |
| Applications: | WB |
| Recommended Dilution: | WB: 0.5-2ug/ml. |
| Reactivity: | Human (Expected from sequence similarity: Mouse, Rat, Dog, Cow) |
| Host: | Goat |
| Isotype: | IgG |
| Clonality: | Polyclonal |
| Immunogen: | Peptide with sequence C-EKDSPRLTKTNP, from the internal region of the protein sequence according to NP_003480.2. |
| Formulation: | 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin |
| Purification: | Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide. Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Aliquot and store at -20C. Minimize freezing and thawing. |
| Conjugation: | Unconjugated |
| Storage: | Store at -20°C as received. |
| Stability: | Stable for 12 months from date of receipt. |
| Gene Name: | nuclear receptor interacting protein 1 |
| Database Link: | NP_003480 Entrez Gene 268903 Mouse Entrez Gene 304157 Rat Entrez Gene 478385 Dog Entrez Gene 8204 Human P48552 |
| Background: | Nuclear receptor interacting protein 1 (NRIP1) is a nuclear protein that specifically interacts with the hormone-dependent activation domain AF2 of nuclear receptors. Also known as RIP140, this protein modulates transcriptional activity of the estrogen receptor. [provided by RefSeq] |
| Synonyms: | RIP140 |
| Protein Families: | Druggable Genome, Transcription Factors |



[View online »](#)

Product images:

TA305768 (0.5ug/ml) staining of Human Cerebellum lysate (35ug protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.