

## **Product datasheet for TA305999**

## 9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US

OriGene Technologies, Inc.

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## **MYD88 Rabbit Polyclonal Antibody**

**Product data:** 

**Product Type:** Primary Antibodies

**Applications:** ELISA, IF, IP, WB

**Recommended Dilution:** WB: 2 μg/mL; IF: 20 μg/mL.

Antibody validated: Western Blot in human, mouse and rat samples; Immunofluorescence and Immunoprecipitation in human samples. All other applications and species not yet

tested.

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: MyD88 antibody was raised against a peptide corresponding to amino acids 279 to 296 of

human MyD88, which are identical to those of mouse.

**Formulation:** PBS containing 0.02% sodium azide.

Concentration: 1ug/ul

**Purification:** MYD88 Antibody is affinity chromatography purified via peptide column.

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Gene Name: myeloid differentiation primary response 88

Database Link: NP 001166037

Entrez Gene 17874 MouseEntrez Gene 301059 RatEntrez Gene 4615 Human

Q99836





Background:

The pro-inflammatory cytokine IL-1 induced cellular response requires IL-1 receptor complex including IL-1RI and IL-1RAcP. Recently, MyD88 was identified as an adapter molecule in the IL-1 signaling pathway (1). MyD88 associates with and recruits IRAK to the IL-1 receptor complex in response to IL-1 treatment and dominant negative form of MyD88 attenuates IL-1R-mediated NF-kappaB activation. MyD88 is also employed as a regulator molecule by IL-18 receptor and human Toll receptor (2,3), which are members in the Toll/IL-1R family of receptors. Targeted disruption of the MyD88 gene results in lose of cellular responses to IL-1 and IL-18, and MyD88-deficient mice lack responses to bacterial product LPS that employs Toll-like receptors 2 and 4 (TLR2 and TLR4) as the signaling receptors. MyD88 is a general adapter protein for the Toll/IL-1R family of receptors and plays an important role in the inflammatory response induced by cytokines IL-1 and IL-18 and endotoxin. MyD88 gene is expressed in many tissues.

Synonyms: MYD88D

**Protein Families:** Druggable Genome

**Protein Pathways:** Apoptosis, Toll-like receptor signaling pathway