

## **Product datasheet for TA306022**

## 9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

OriGene Technologies, Inc.

## **APRIL (TNFSF13) Rabbit Polyclonal Antibody**

**Product data:** 

**Product Type:** Primary Antibodies

Applications: WB

Recommended Dilution: WB: 5 - 10 ug/mL

Reactivity: Human

Host: Rabbit

Isotype: IgG

Clonality: Polyclonal

**Immunogen:** APRIL antibody was raised against a synthetic peptide corresponding to amino acids near the

center of human APRIL.

**Formulation:** PBS containing 0.02% sodium azide.

**Concentration:** 1ug/ul

**Purification:** Affinity chromatography purified via peptide column

**Conjugation:** Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Gene Name:** tumor necrosis factor superfamily member 13

Database Link: NP 003799

Entrez Gene 8741 Human

<u>O75888</u>

**Background:** Members in the TNF superfamily regulate immune responses and induce apoptosis. A novel

member in the TNF family was recently identified by several groups and designated APRIL (for a proliferation-inducing ligand), TALL-2 (for TNF- and ApoL-related Leukocyte-expressed Ligand 2), and TRDL-1alpha (for TNF related death ligand 1a) in human and mouse (1-4). Two receptors for APRIL were recently identified and designated TACI and BCMA (5-7). APRIL stimulates B and T cell proliferation, triggers humoral immune responses, activates NF-kappaB, and induces cell death (1-7). APRIL and its close relative of BlyS and their receptors

BCMA and TACI are involved in diseases of autoimmunity and cancer (8,9).

Synonyms: APRIL; CD256; PRO715; TALL-2; TALL2; TNLG7B; TRDL-1; UNQ383; ZTNF2






**Protein Families:** Druggable Genome, Secreted Protein, Transmembrane

**Protein Pathways:** Cytokine-cytokine receptor interaction

## **Product images:**



Western blot analysis of APRIL expression in K562 cells with APRIL antibody at (A) 5 and (B) 10 ug/ml.