

Product datasheet for **TA306038**

AIF (AIFM1) Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	IHC, WB
Recommended Dilution:	WB: 1 ug/mL, ICC: 10 ug/mL
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Isotype:	IgG
Clonality:	Polyclonal
Immunogen:	AIF antibody was raised against a peptide corresponding to amino acids 517 to 531 of human AIF. This sequence is identical to those of mouse and rat AIF.
Formulation:	PBS containing 0.02% sodium azide.
Concentration:	1ug/ul
Purification:	Ion exchange chromatography purified
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Gene Name:	apoptosis inducing factor, mitochondria associated 1
Database Link:	NP_001124318 Entrez Gene 26926 Mouse Entrez Gene 83533 Rat Entrez Gene 9131 Human O95831



[View online »](#)

Background:

Apoptosis is characterized by several morphological nuclear changes including chromatin condensation and nuclear fragmentation. These changes are triggered by the activation of members of caspase family, caspase activated DNase, and several novel proteins (1). A novel gene, the product of which causes chromatin condensation and DNA fragmentation, was recently identified, cloned, and designated apoptosis inducing factor (AIF) (2). Like the critical molecules, cytochrome c and caspase-9, in apoptosis, AIF localizes in mitochondria. AIF translocates to the nucleus when apoptosis is induced and induces mitochondria to release the apoptogenic proteins cytochrome c and caspase-9. AIF induces chromatin condensation and large scale DNA fragmentation, which are the hallmarks of apoptosis, of the isolated nucleus and the nucleus in live cells by microinjection and apoptosis stimuli (2,3). AIF is highly conserved between human and mouse and widely expressed (2).

Synonyms:


AIF; CMT2D; CMTX4; COWCK; COXPD6; NADMR; NAMSD; PDCD8

Protein Families:


Druggable Genome, Transmembrane

Protein Pathways:

Apoptosis

Product images:


Western blot analysis of AIF in K562 cell lysate (A), rat heart (B), and mouse heart (C) tissue lysates with AIF antibody (IN) at 1 ug/ml.



Immunohistochemistry of AIF in human retina with AIF antibody at 10 ug/ml.