

## Product datasheet for **TA306145**

### XAF1 Rabbit Polyclonal Antibody

#### Product data:


Product Type:	Primary Antibodies
Applications:	IF, IHC, WB
Recommended Dilution:	WB: 0.5 - 2 ug/mL, ICC: 2 ug/mL, IF: 10 ug/mL
Reactivity:	Human, Mouse
Host:	Rabbit
Isotype:	IgG
Clonality:	Polyclonal
Immunogen:	XAF-1 antibody was raised against a synthetic peptide corresponding to amino acids at the C-terminus of human XAF-1.
Formulation:	PBS containing 0.02% sodium azide.
Concentration:	1ug/ul
Purification:	Affinity chromatography purified via peptide column
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Gene Name:	XIAP associated factor 1
Database Link:	<a href="#">CAA68030</a> <a href="#">Entrez Gene 54739 Human</a> <a href="#">Q6GPH4</a>
Background:	XAF-1 binds to XIAP, an inhibitor of caspases-3, -7, and -9, and triggers its relocation from the cytosol to the nucleus (1,2). Overexpression of XAF-1 results in the neutralization of XIAP's ability to inhibit cell death (1). XAF-1 is normally expressed in all adult and fetal tissues but was found to be present in very low levels in a variety of cancer cell lines (3). In contrast, XIAP levels have been shown to be high in a majority of cell lines. Low XAF-1 and high basal expression of XIAP may therefore play a critical role in maintaining survival of cancer cell lines (2,3). Both IFN-alpha2 and IFN-beta can induce XAF-1 mRNA in all cells examined but induction of XAF-1 protein (as observed by immunoblot analysis) was seen only in cell lines sensitive to the apoptotic effects of IFNs (4).




[View online »](#)

Synonyms: BIRC4BP; HSXIAPAF1; XIAPAF1


### Product images:



Western blot analysis of XAF-1 in human spleen lysate with XAF-1 antibody at 0.5 (lane A), 1 (lane B), and 2 (lane C) ug/ml, respectively.



Immunohistochemical staining of human spleen tissue using XAF-1 antibody at 2 ug/ml.



Immunofluorescence of XAF-1 in Human Spleen tissue with XAF-1 antibody at 10 ug/mL.