

Product datasheet for TA306208

MCL1 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IF, WB

Recommended Dilution: WB: 1 - 2 ug/mL, ICC: 10 ug/mL, IF: 20 ug/mL

Reactivity: Human
Host: Rabbit
Isotype: IgG

Clonality: Polyclonal

Immunogen: Mcl-1 antibody was raised against a peptide corresponding to 16 amino acids near the

amino-terminus of human Mcl-1.

Formulation: PBS containing 0.02% sodium azide.

Concentration: 1ug/ul

Purification: Ion exchange chromatography purified

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Gene Name: myeloid cell leukemia 1

Database Link: NP 068779

Entrez Gene 4170 Human

Q07820

Background: Myeloid cell leukimia-1 (Mcl-1) is a member of the Bcl-2 family of proteins that can act to

promote cell survival (reviewed in 1). While the mechanism by which Mcl-1 inhibits apoptosis is not known, it is thought that it may heterodimerize and neutralize pro-apoptotic members of the Bcl-2 family such as Bim or Bak (2,3). Mcl-1 was originally identified in differentiating myeloid cells (4), but has since been shown to be expressed in multiple cell types. Mcl-1 is essential for embryogenesis (5) and for the development and maintenance of B and T lymphocytes in animals (3). Mcl-1 exists as at least two distinct isoforms designated Mcl-1L and Mcl-1S. In marked contrast to the larger isoform of Mcl-1, overexpression of Mcl-1S

promotes cell death (6).



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Synonyms: bcl2-L-3; BCL2L3; EAT; Mcl-1; mcl1; MCL1-ES; MCL1L; MCL1S; TM

Protein Families: Druggable Genome, Transmembrane


Product images:



Western blot analysis of Mcl-1in Raji cell lysates with Mcl-1 antibody at (A) 1 and (B) 2 ug/mL.



Immunocytochemistry of McI-1 in Raji cells with McI-1 antibody at 10 ug/mL.



Immunofluorescence of Mcl-1 in Raji cells with Mcl-1 antibody at 20 ug/mL.